

# Using EOS Imaging Technique to Provide Reference Values for Proximal, Distal, and Total Thoracic Kyphosis in the Iranian Population

Ghandehari Hasan<sup>1</sup>, Tanhapour Samaneh<sup>2</sup>, Hassanzadeh Mohammad<sup>3</sup>, Rooyan Pooria<sup>1</sup>✉

1. Department of Orthopedics, Iran University of Medical Sciences, Tehran, Iran

2. Student Research Committee, Iran University of Medical Sciences, Tehran, Iran

3. Bone and Joint Reconstruction Research Center, Department of Orthopedics, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

## Article Info

### Article Type:

Original Article

### Article History:

Received

23 Oct 2022

Received in revised form

09 Nov 2022

Accepted

02 Jan 2023

Published online

01 Mar 2023

### Publisher:

Fasa University of  
Medical Sciences

## Abstract

**Background & Objectives:** EOS, as a new imaging technique, has several benefits over conventional radiography in measuring spinal curvatures. In this study, we aimed to use EOS to determine the normal values of thoracic kyphosis besides its association with the participants' age and gender.

**Materials & Methods:** In this retrospective cross-sectional study, 100 individuals with no history of spinal abnormality were included. Spinal curvatures, comprising proximal thoracic segment, distal thoracic segment, and total thoracic kyphosis, were measured on EOS images using a Cobb method. The demographic characteristics of the participants, such as age and gender were also extracted from medical profiles.

**Results:** The study population included 31 males and 69 females with a mean age of  $27.2 \pm 17.4$  years (range 6-66). The mean curvatures of proximal and distal thoracic segments were  $14.5 \pm 6.5^\circ$  and  $32.7 \pm 7.5^\circ$ , respectively. The mean total thoracic kyphosis was  $47.2 \pm 7.2^\circ$ . Besides, distal and total thoracic kyphosis, but not proximal thoracic segment, were significantly more in participants aged over 40 years ( $p=0.03$ ,  $p=0.04$ , and  $p=0.07$ , respectively). No significant association was found between the thoracic curvatures and participants' sex.

**Conclusion:** It seems that there is a direct relationship between distal and total thoracic kyphosis and aging; however, more studies are required in order to approve our results.

**Keywords:** Spinal curvature, thoracic kyphosis, EOS

**Cite this article:** Ghandehari H, Tanhapour S, Hassanzadeh M, Rooyan P. Using EOS Imaging Technique to Provide Reference Values for Proximal, Distal, and Total Thoracic Kyphosis in the Iranian Population. JABS. 2023; 13 (1): 23-28.

**DOI:** 10.18502/jabs.v13i1.12074

## Introduction

In a normal human, the thoracic spine contains some degree of kyphosis. This kyphosis can be divided into two segments, including proximal thoracic kyphosis (from T1 to T5) and distal thoracic kyphosis (from T5 to T12).

Each segment has a specific effect on the total thoracic curvature of the spine (1).

Thoracic kyphosis is affected by several physiologic conditions such as aging, and pathologic conditions such as fracture, osteoarthritis, and congenital anomalies (2).

While the angle of kyphosis is more than normal ranges, hyperkyphosis occurs, which is generally assumed as a Cobb angle

✉ **Corresponding Author: Rooyan Pooria**, Department of Orthopedics, Iran University of Medical Sciences, Tehran, Iran  
Email: [rooyanpooria@gmail.com](mailto:rooyanpooria@gmail.com)

of 50° or more in the standing position and has several detrimental effects through changing vertebral loading during daily activities (3, 4). Therefore, reliable evaluation of thoracic kyphosis has critical importance.

To date, a standing lateral spine X-ray is the gold standard for the objective assessment of kyphosis (5). However, the reliability of kyphosis measurement on lateral spine radiographs is small (6). Therefore, there are strong attempts to find a safe and reliable method for kyphosis measurement. EOS is a 3D imaging system with low-dose radiation capable of capturing full-body 3D images of patients in a natural and standing position (6). It is shown that patients with adolescent idiopathic scoliosis will receive approximately 16–34 times lesser organ dose from EOS as compared with the standard digital radiography (6). Moreover, it produces images with higher contrast and sharpness; thereby, it can provide more reliable and accurate measurement of spinal curvatures (7). For these advantages, there is a growing trend towards the implication of EOS for measuring spinal curvatures (8, 9).

Recently, several studies have investigated the implication of EOS imaging for the evaluation of thoracic kyphosis (8, 10, 11). However, according to our knowledge, no study has been performed to define the normal value of the proximal thoracic segment using EOS. Hence, in this work, we aimed to harness the EOS imaging technique for the assessment of the normal values

for thoracic kyphotic parameters, such as spinal curvatures of proximal and distal segments, in the asymptomatic Iranian population. We also assessed the association of the measured angle with the participants' age and gender.

## Materials & Methods

This retrospective cross-sectional study was approved by the Ethics Committee of Iran University of Medical Sciences (ethical code: IR.IUMS.FMD.REC.1398.255). Medical profiles of the individuals who underwent EOS imaging in the Shafa Orthopedic hospital between January 2019 and January 2020 were reviewed. Individuals with a history of spinal abnormality were excluded from the study. The remaining patients were included in the study (n=100).

The demographic characteristics of the participants, including age and gender, were extracted from the participants' medical records. Spinal curvatures were extracted from the EOS images and included proximal thoracic segment, distal thoracic segment, total thoracic kyphosis, and lumbar lordosis (Figure 1). The Cobb method was used to measure the proximal (sagittal angle between T1 and T5), distal (sagittal angle between T5 and T12), and total thoracic kyphosis angle (angle between T1 and T12). Lumbar lordosis was assessed through the Cobb method between the line along with the superior L1 end plate and the line along with the S1 endplate.



**Figure 1.** Evaluation of total kyphosis and lumbar lordosis on the EOS image

## EOS Imaging Technique for Thoracic Kyphosis

## Statistical Analysis

SPSS software, version 16, was used for the statistical analysis. Descriptive statistics were provided with mean  $\pm$  standard deviation or number & percentage. A comparison of mean values between different groups was made by an independent t-test or its non-parametric counterpart (Mann-Whitney U test). P values less than 0.05 were considered statistically significant.

## Results

A total of 100 participants with no history of spinal abnormality were included in this study. The study population included 31 males and 69 females with a mean age of  $27.2 \pm 17.4$  years (range 6-66). The most

common age group was 10-20 years ( $n=44$ ).

The mean proximal thoracic segment was  $14.5 \pm 6.5^\circ$ . The mean distal thoracic segment was  $32.7 \pm 7.5^\circ$ . The mean total thoracic kyphosis was  $47.2 \pm 7.2^\circ$ . The mean lumbar lordosis was  $57.6 \pm 11.1^\circ$ . Spinal curvatures were not significantly different between males and females (Table 1). Distal and total thoracic kyphosis was significantly more in participants aged over 40 years. Proximal thoracic kyphosis was not significantly different between the participants aged  $<40$  years and participants aged  $>40$  years. Lumbar lordosis was significantly smaller in participants older than 40 years of age in comparison with those younger than 40 years of age (Table 2).

**Table 1.** Comparison of spinal curvatures between males and females

Variable	Males (n=31)	Females (n=69)	P-value
Proximal thoracic segment ( $^\circ$ )	$14.6 \pm 6.5$	$14.4 \pm 6.5$	0.88
Distal thoracic segment ( $^\circ$ )	$33.3 \pm 8.1$	$32.3 \pm 7.3$	0.53
Total thoracic kyphosis ( $^\circ$ )	$48 \pm 8.5$	$46.8 \pm 6.6$	0.44
Lumbar lordosis ( $^\circ$ )	$56.8 \pm 11.2$	$58 \pm 11$	0.61

Data are presented as mean  $\pm$  standard deviation. P-value  $<0.05$  is considered significant

**Table 2.** Comparison of spinal curvatures between males and females

Variable	Aged <40 y (n=73)	Aged >40 y (n=27)	P-value
Proximal thoracic segment (°)	14.2±6.4	14.8±6.7	0.07
Distal thoracic segment (°)	32.2±8.6	34.5±8	0.03
Total thoracic kyphosis (°)	46.1±8.3	48.8±6.9	0.04
Lumbar lordosis (°)	58.6±11	56.2±10.8	0.03

Data are presented as mean ± standard deviation. P-value <0.05 is considered significant

## Discussion

Hyperkyphosis is described as abnormal thoracic spine curvature, and the Cobb method is a gold test for its diagnosis (12, 13). If this condition is not rapidly recognized and stopped, serious problems may occur, like osteoporosis. To detect this abnormal condition, we must have enough information about the normal range of these angles (14). On the other hand, EOS is a modern imaging technique based on a slot-scanning X-ray system to diminish radiation exposure in orthopedic-related images and has barely been used for the measurement of spinal curvatures (15).

Hence, in this study, we assessed the normal values of proximal, distal, and total thoracic kyphosis in an Iranian population with no history of spinal abnormality by this

imaging method. Our results indicated that the evaluated variables were not associated with the participants' gender. However, distal and total thoracic kyphosis was significantly more in participants older than 40 years of age.

In another attempt, Gleb et al. determined the indices of sagittal spinal alignment on standing radiographs of 100 adults older than 40 and without a history of significant spinal abnormality (16). In this work, the mean angle of proximal and distal thoracic segments were  $14 \pm 8$  and  $34 \pm 11$ , and unlike our findings (the mean angle of proximal and distal thoracic segments were  $14.8 \pm 6.7$  and  $34.5 \pm 8$ , respectively). No significant association was found between the thoracic kyphosis and the age of the patients (16). This difference may be associated with the exerted inclusion criteria

for selecting subjects. In the study of Gleb et al. only adults aged >40 years were included whereas we had no age limitation. Also, Abrisham and co-workers assessed the normal range of thoracic kyphosis and lumbar lordosis angles in the Iranian population using EOS imaging method. The average angles of thoracic kyphosis and lumbar lordosis were  $43.55 \pm 6.44$  and  $32.42 \pm 6.29$ , respectively, and dissimilar to our reports, the mean lumbar lordosis and thoracic kyphosis were significantly associated with sex but not with age (8). This difference could be attributed to the higher number of participants aged over 40 in the study of Abrisham et al.

In another research by Bassani et al., EOS imaging technique was used to describe the normative parameter of sagittal spine alignment in 160 asymptomatic elderly subjects (age >60, Caucasian). In the mentioned research, the mean angles of thoracic kyphosis and lumbar lordosis were  $55 \pm 14^\circ$  and  $57 \pm 12^\circ$ , and finally, they concluded that the reference values in the elderly are considerably different compared to younger adults (11). Similarly, the total mean thoracic kyphosis and lumbar lordosis was  $48.8 \pm 6.9^\circ$  and  $56.2 \pm 10.8^\circ$  in participants aged > 40 years. In the current study that was significantly different from what was observed in adults aged <40 years. It has been declared that the aging occurrence changes postural alignment in light of morphostructural alteration of the elements involved in the stability of posture (17).

The main limitations were retrospective design, lack of a control group assessed by conventional radiography, a small number of patients in some age subgroups, and inability to perform multivariate analysis. Therefore, future complementary studies are required to confirm the results of this study.

### **Conclusions**

Our study showed that there is a direct relationship between distal and total thoracic kyphosis and aging. Also, thoracic kyphosis

angle could be assessed by the EOS imaging method, allowing a lesser organ irradiation dose and sharpen images. However, more projects are needed in order to validate our results.

### **Data Availability Statement**

All data related to this research have been included in the manuscript.

### **Acknowledgement**

The present study was supported by Iran University of Medical Sciences (ethic code: IR.IUMS.FMD.REC.1398.255).

### **Conflict of Interest**

The authors have no conflicts of interest to declare.

### **References**

1. Moreira Pinto E, Alves J, de Castro AM, Silva M, Miradouro J, Teixeira A, et al. High thoracic kyphosis: impact on total thoracic kyphosis and cervical alignment in patients with adolescent idiopathic scoliosis. *Spine deformity*. 2020 Aug;8:647-53.
2. Perriman DM, Scarvell JM, Hughes AR, Lueck CJ, Dear KB, Smith PN. Thoracic hyperkyphosis: a survey of Australian physiotherapists. *Physiotherapy Research International*. 2012 Sep;17(3):167-78.
3. Bruno AG, Anderson DE, D'Agostino J, Boussein ML. The effect of thoracic kyphosis and sagittal plane alignment on vertebral compressive loading. *Journal of Bone and Mineral Research*. 2012 Oct;27(10):2144-51.
4. Koelé MC, Lems WF, Willems HC. The clinical relevance of hyperkyphosis: a narrative review. *Frontiers in endocrinology*. 2020;11:5.
5. Katzman WB, Wanek L, Shepherd JA, Sellmeyer DE. Age-related hyperkyphosis: its causes, consequences, and management. *Journal of orthopaedic & sports physical therapy*. 2010;40(6):352-60.
6. Kuklo TR, Potter BK, Polly Jr DW, O'Brien MF, Schroeder TM, Lenke LG. Reliability analysis for manual adolescent idiopathic scoliosis measurements. *Spine*. 2005;30(4):444-54.
7. Kim SB, Heo YM, Hwang CM, Kim TG, Hong JY, Won YG, et al. Reliability of the EOS Imaging System for Assessment of the Spinal and Pelvic Alignment in the Sagittal Plane. *Clin Orthop Surg*. 2018;10(4):500-7.
8. Abrisham SM, Ardekani MR, Mzarch MA. Evaluation of the normal range of thoracic kyphosis and lumbar lordosis

- angles using EOS imaging. *Maedica*. 2020;15(1):87.
9. Dubousset J, Ilharreborde B, Le Huec JC. Use of EOS imaging for the assessment of scoliosis deformities: application to postoperative 3D quantitative analysis of the trunk. *European Spine Journal*. 2014;23:397-405.
  10. Le Huec J, Demezou H, Aunoble S. Sagittal parameters of global cervical balance using EOS imaging: normative values from a prospective cohort of asymptomatic volunteers. *European Spine Journal*. 2015;24(1):63-71.
  11. Bassani T, Galbusera F, Luca A, Lovi A, Gallazzi E, Brayda-Bruno M. Physiological variations in the sagittal spine alignment in an asymptomatic elderly population. *The Spine Journal*. 2019;19(11):1840-9.
  12. Fukuoka Y, Katzman WB, Gladin A, Lane NE, Kado DM, Oh YJ. Slower upper extremity function in older adults with hyperkyphosis negatively impacts the 6-min walk test. *BMC Musculoskeletal Disorders*. 2022;23(1):505.
  13. Kado DM, Miller-Martinez D, Lui LY, Cawthon P, Katzman WB, Hillier TA, et al. Hyperkyphosis, kyphosis progression, and risk of non-spine fractures in older community dwelling women: the Study of Osteoporotic Fractures (SOF). *Journal of Bone and Mineral Research*. 2014;29(10):2210-6.
  14. Abrisham SMJ, Ardekani MRS, Mzarch MAB. Evaluation of the Normal Range of Thoracic Kyphosis and Lumbar Lordosis Angles Using EOS Imaging. *Maedica (Bucur)*. 2020;15(1):87-91.
  15. Luo TD, Stans AA, Schueler BA, Larson AN. Cumulative Radiation Exposure With EOS Imaging Compared With Standard Spine Radiographs. *Spine Deformity*. 2015;3(2):144-50.
  16. Gelb DE, Lenke LG, Bridwell KH, Blanke K, McEneaney KW. An analysis of sagittal spinal alignment in 100 asymptomatic middle and older aged volunteers. *Spine*. 1995;20(12):1351-8.
  17. Regolin F, Carvalho GA. Relationship between thoracic kyphosis, bone mineral density, and postural control in elderly women. *Brazilian Journal of Physical Therapy*. 2010;14:464-9.