




## Comparison of Treatment Outcomes for Tennis Elbow: Corticosteroid Injection and Casting vs. Acupuncture and Physiotherapy

Heydari Sina<sup>1</sup>, Jafarnezhad Aboubakr<sup>2</sup>, Faryabi Reza<sup>3</sup>, Raesi Rasoul<sup>4,5</sup>, Goroohi Mahya<sup>6</sup>  
Mehralizade Akbar<sup>3</sup>, Amiri Ghanatsaman Fatemeh<sup>7</sup>, Daneshi Salman<sup>3</sup>✉

1. School of Medicine, Imam Khomeini Hospital, Jiroft University of Medical Science, Jiroft, Iran
2. Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran
3. Department of Public Health, School of Health, Jiroft University of Medical Sciences, Jiroft, Iran
4. Department of Nursing, Torbat Jam Faculty of Medical Sciences, Torbat Jam, Iran
5. Department of Health Services Management, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran
6. Clinical Research Development Center, Imam Khomeini Hospital, Jiroft University of Medical Sciences, Jiroft, Iran
7. Student Research Committee, Faculty of Nursing and Midwifery, Kerman University of Medical Science, Kerman, Iran

### Article Info

#### Article Type:

Original Article

#### Article History:

Received

12 Dec 2023

Received in revised form

30 Dec 2023

Accepted

01 Jan 2024

Published online

15 Jan 2024

#### Publisher:

Fasa University of  
Medical Sciences

### Abstract

**Background & Objectives:** Today, despite the existence of numerous treatments for tennis elbow, the search for more effective methods continues due to the limited efficacy of these existing approaches. This study aimed to compare the results of tennis elbow treatment in two groups: patients receiving corticosteroid injection and casting, and patients receiving acupuncture and physiotherapy.

**Materials & Methods:** This quasi-experimental study, conducted in Jiroft in 2021, investigated the effectiveness of different treatment approaches for tennis elbow. Fifty patients diagnosed with tennis elbow were randomly divided into two groups of 25 each. One group received a combination of corticosteroids and casts, while the other group received acupuncture and physiotherapy. A pain line was used to assess pain levels based on VAS criteria. Data were analyzed using SPSS-26 statistical software, employing Chi-square, Fisher's Exact, Mann-Whitney, one-way ANOVA, Kruskal-Wallis, and Kolmogorov-Smirnov tests.

**Results:** The majority of patients were female (66%) and housewives (46%). There was a significant difference in pain outcomes between the two groups: physiotherapy alone and physiotherapy combined with corticosteroids and casts. This difference was observed across various conditions (pain at rest, pain during activity, and amount of pain evoked during activity) ( $p < 0.05$ ). In simpler terms, corticosteroids and casts were more effective in reducing pain from tennis elbow compared to acupuncture and physiotherapy. Additionally, corticosteroids and casts had a greater effect on hand movement (supination, pronation, extension, and flexion) compared to acupuncture and physiotherapy.

**Conclusion:** Our findings indicate that treatment of tennis elbow with a combination of corticosteroid injection and casting is more effective than acupuncture or physiotherapy alone. Corticosteroid injections themselves are also an effective way to relieve tennis elbow pain.

**Keywords:** Tennis Elbow, therapeutics, injections, acupuncture, orthopedics, patients

**Cite this article:** Heydari S, Jafarnezhad A, Faryabi R, Raesi R, Goroohi M, Mehralizade A, Amiri Ghanatsaman F, Daneshi S. Comparison of Treatment Outcomes for Tennis Elbow: Corticosteroid Injection and Casting vs. Acupuncture and Physiotherapy. JABS.2024; 14(1): 67-76.

**DOI:** 10.18502/jabs.v14i1.14808

### Introduction

Tennis elbow is a common soft tissue injury of the elbow caused by overuse and micro-tears in the extensor muscles that extend the wrist. While frequently seen in athletes, it can affect anyone who performs repetitive motions in their daily activities or jobs. Professions such as painting,

butchery, carpentry, and plumbing carry a higher risk due to the repetitive movements required (1). Tennis elbow presents as pain and tenderness on the outer elbow, sometimes radiating to the forearm and back of the hand (2-4). These symptoms often worsen with arm use and frequent wrist movements (5). The condition is most common between the ages of 35 and 55 and typically affects the dominant arm (6-8). Recovery time for tennis elbow can range from 6 months to 2 years (9).

✉ **Corresponding Authors:** Daneshi Salman, Department of Public Health, School of Health, Jiroft University of Medical Sciences, Jiroft, Iran  
Email: [salmandaneshi008@gmail.com](mailto:salmandaneshi008@gmail.com)



The most common causes of tennis elbow are overuse and repetitive straining of the forearm muscles and tendons. However, a direct blow or collision to the elbow can also cause it (10-12). Diagnosis usually relies on a medical history and physical examination, though imaging tests may be needed in some cases (13).

Studies have shown that the wrist extensor carpi radialis brevis tendon is the most common source of pain and dysfunction in patients with tennis elbow (14). However, no single, specific therapeutic intervention has been identified as the most effective treatment (15). Tennis elbow can heal on its own, but this process may take several weeks or months (16, 17). Simple treatments such as rest, cold compresses (plural), and avoiding activities that aggravate the injured tendons and muscles can alleviate the pain (18, 19). Invasive treatments such as surgery are reserved only for severe and resistant cases (20). Individuals with jobs that involve frequent manual tasks, such as lifting objects, should avoid such activities until the pain subsides (21).

Several non-invasive treatments have been proposed for tennis elbow (22), including exercise therapy (23-25), corticosteroid injections (26, 27), medication (28, 29), laser therapy (30), electrical stimulation (31, 32), ergonomic modifications (33), bracing with counterforce, acupuncture (34, 35), and splinting. While corticosteroid injections are common for persistent tennis elbow (36), they can have negative side effects and high recurrence rates (37). Similarly, the effectiveness of acupuncture and dry needling requires further investigation (35). Physiotherapy and short-term use of orthotics have shown promise in reducing pain and improving blood flow (38). Surgery remains a last resort for severe and chronic cases (39). Currently, no single treatment demonstrates definitive superiority, highlighting the need for further research to identify the most effective approach for tennis elbow (40).

This has resulted in many beta errors in studies, thereby significantly reducing their ability to detect group differences. Consequently, the present study compares two treatment methods for tennis elbow: corticosteroid injection and casting versus acupuncture and physiotherapy.

## Materials & Methods

The present study is quasi-experimental. The study population consisted of patients with tennis elbow who were referred to orthopedic clinics based on the diagnoses of specialist physicians. Inclusion criteria were pain and tenderness around the lateral epicondyle, which worsened with active wrist extension, a positive result on one of the diagnostic tests (Kazen 3 or Mills 4), and pain when making a fist.

Exclusion criteria also included injections into the affected area within the previous 6 months of a diagnosis of diabetes, pregnancy, peripheral nerve entrapment at the doctor's discretion, and cervical radiculopathy. Participants who were eligible provided informed consent. They completed and signed the consent form to participate in the research project. Then, all participants participated in the pretest to evaluate the research variables including pain at rest, pain during activity, and evoked pain.

In the present study, a total of 40 patients were evaluated based on previous studies (41). Twenty patients received corticosteroids and limb plastering, while another twenty received acupuncture and physiotherapy. The sample size was calculated with a 95% confidence level and 90% power based on the results of pilot studies involving 20 people. It was calculated using the following formula so that 25 people were allocated to each group, resulting in a total sample size of 50 people across the groups of corticosteroid and limb plaster recipients, as well as acupuncture and physiotherapy recipients.

$$n = \frac{(s_1^2 + s_2^2) \left( z_{1-\alpha} + z_{1-\beta} \right)^2}{d^2} = \frac{(s_1^2 + s_2^2) \left( z_{1-\alpha} + z_{1-\beta} \right)^2}{(\bar{x}_2 - \bar{x}_1)^2}$$

$$= \frac{(5.85 + 6.01)(1.64 + 1.56)^2}{(4.5 - 2.94)^2}$$

$$\cong 50$$

**d2:** is the difference between the means of the two groups, which has reached the power of two

In this study, a pain ruler was used to determine the amount of pain based on VAS criteria. analogical measure of vision is linear pain 100 mm long,

#### Comparison of Corticosteroids with Acupuncture

with one end of the line indicating maximum pain (score 100) and the other end indicating no pain.

To measure pain, the person was instructed to determine the severity of the pain according to the length of the line. The validity and reliability of this device in measuring pain were very high. In this blinding method, the person marked the location of pain on the line. Then, by placing a graduated ruler on the line, the researcher measured the amount of pain in millimeters. To assess aroused pain, the subjects were asked to rate their pain specifically during the chin test, when the hand had no activity. To assess pain at rest, participants were asked to rate their pain when the hand was inactive. To assess pain during activity, participants were asked to rate their pain during an activity they performed in the past 24 hours. Following the pre-test, the participants were randomly divided into two groups: one receiving a corticosteroid injection and plastering, and the other receiving acupuncture and physiotherapy. Data collection involved demographic information questionnaires and follow-up information based on the researcher's checklist and VAS test. After data collection and initial processing, the data were entered into SPSS statistical software version 26 for descriptive and inferential analysis. To address the research questions and objectives, descriptive statistical

methods were used first. These included one- and two-dimensional frequency distribution tables, statistical graphs, and descriptive statistics indices. Inferential statistical tests were then employed, including Fisher's exact test, chi-square ( $X^2$ ) test, Student's t-tests, and one-way analysis of variance. All participants in this research provided informed consent and participated voluntarily. Before conducting surveys and measurements, the researcher provided participants with all necessary information about the study. Participants were also reassured about the anonymity and confidentiality of their data. Data collection adhered to all relevant rules and regulations. This study builds upon a General Medicine dissertation conducted at Jiroft University of Medical Sciences.

#### Results

This study involved 50 participants: 25 received corticosteroids and plastering, while the other 25 received acupuncture and physiotherapy. Table 1 presents the mean age, standard deviation, minimum and maximum values, and the results of the two-independent-samples t-test used to compare the age of the acupuncture and physiotherapy group with the corticosteroid and plastering group.

**Table 1.** Results of independent t-test to compare the two groups of acupuncture and physiotherapy and corticosteroids and casts in terms of age

Groups	Minimum (maximum)	Mean $\pm$ SD	P-value
Acupuncture and physiotherapy	29 (58)	44.4 $\pm$ 8.63	0.79
Corticosteroids and casts	25 (66)	43.72 $\pm$ 9.28	

The mean age in the acupuncture and physiotherapy group was 44.48 years (SD = 6.63), while the mean age in the corticosteroid and plastering group was 43.72 years. The age range in the acupuncture and physiotherapy group was 29 to 58 years, while the range in the corticosteroid and plastering group was

25 to 66 years. Although the acupuncture and physiotherapy group appeared to have a slightly higher average age, this difference was not statistically significant ( $p > 0.05$ ).

According to Table 2, 28% of participants in the acupuncture and physiotherapy group were men, while 40% of participants in the

corticosteroid and plastering group were men. Overall, the study population was 34% male (17 people) and 66% female (33 people). There was no statistically significant difference in gender distribution between the two groups ( $p > 0.05$ ).

Occupationally, housewives formed the majority in both the acupuncture and physiotherapy group and the corticosteroid and plastering group. No statistically significant difference ( $p > 0.05$ ) was found in occupational distribution between the groups.

**Table 2.** Frequency distribution of sex and job in treatment groups by acupuncture, physiotherapy, corticosteroids, and casts

Variable	Groups		P-Value
	Acupuncture and physiotherapy	Corticosteroids and casts	
Male	7 (28.00)	10 (40.00)	0.22
Female	18 (72.00)	15 (60.00)	
Job			
Employee	6 (24.00)	2 (8.00)	
Self-employment	3 (12.00)	2 (8.00)	
Farmer	4 (16.00)	9 (36.00)	
Unemployed	0 (0.00)	1 (4.00)	
Housewife	12 (48.00)	11 (44.00)	

Table 3 presents the results of the t-tests comparing the acupuncture and physiotherapy group with the corticosteroid and plastering group on four measures: pain at rest, pain during activity, amount of pain evoked during activity, and weight gain in kilograms. The acupuncture and physiotherapy group appeared to have

a higher average pain score at rest compared to the corticosteroid and plastering group. This difference was statistically significant ( $p < 0.05$ ). Conversely, the corticosteroid and plastering group experienced a statistically significant ( $p < 0.05$ ) increase in weight gain compared to the acupuncture and physiotherapy group.

**Table 3.** The T-test to compare the two groups of treatment with acupuncture, physiotherapy, corticosteroids, and casts in terms of pain at rest, pain during activity, amount of pain evoked during activity, and weight gain in kilograms

Variable	Treatment group		p-value
	Acupuncture and physiotherapy	Corticosteroids and casts	
The amount of pain at rest	6.78±1.84	2.94±2.16	0.001
The amount of pain during activity	5.3±1.42	1.44±1.64	0.001
The amount of pain evoked during activity	4.5±1.56	1.8±2.16	0.001
Weight gain	4.36±2.34	8.08±5.36	0.003

#### Comparison of Corticosteroids with Acupuncture

Table 4 shows the mean rankings and the results of the Mann-Whitney U test used to compare the acupuncture and physiotherapy group with the corticosteroid and plastering group on four movement measures: hand supination rate, hand pronation rate, hand extension rate, and hand flexion rate.

Individuals in the corticosteroid and plastering group received higher average rankings for all four movement measures compared to the acupuncture and physiotherapy group. This difference in treatment methods regarding the studied factors was statistically significant ( $p < 0.05$ ).

**Table 4.** Mann-Whitney test results to compare the two groups of acupuncture and physiotherapy and corticosteroids and plaster casts in terms of the degree of supination of the hand, hand pronation, hand extension, flexion in the hand

Variable	Treatment group	Average ratings	Mann-Whitney U	p-value
The extent of hand supination	Acupuncture and physiotherapy	18.34	133.5	0.001
	Corticosteroids and casts	32.66		
Hand pronation rate	Acupuncture and physiotherapy	18.34	133.5	0.001
	Corticosteroids and casts	32.66		
The number of hand extensions	Acupuncture and physiotherapy	20.9	197.5	0.016
	Corticosteroids and casts	30.1		
Flexion rate in hand	Acupuncture and physiotherapy	19.56	164	0.002
	Corticosteroids and casts	31.44		

## Discussion

Despite the numerous methods proposed for treating tennis elbow in recent decades, insufficient scientific evidence remains to definitively select the best approach (32, 36). This lack of clarity could stem from various factors, including the self-limiting nature of the disease itself (36, 42, 43). Therefore, this study aimed to determine and compare the outcomes of tennis elbow treatment in two groups: patients receiving corticosteroid injection and casting, and patients treated with acupuncture and physiotherapy. The study population primarily consisted of middle-aged females who were housewives. Kivi et al. (44) suggest a possible explanation for the higher prevalence of lateral epicondylitis in women aged 40-60: a decrease in tendon elasticity, particularly at the junction point, coinciding with repetitive biomechanical stress at midlife. Hutson's study (45) reported a similar age range for peak

prevalence (30-55 years old), with a rarity in individuals under 30 and a higher incidence with increasing age. Rayan et al. (46) further support the link between gender and prevalence, noting a higher rate in women and a correlation with age and time spent on hand-intensive tasks. Housewives, in particular, may be more susceptible due to the repetitive nature of their daily activities (gardening, cleaning, cooking) that strain the forearm muscles and tendons.

This study found a significant difference between the acupuncture and physiotherapy group and the corticosteroid and plastering group in terms of pain response across various conditions (pain during rest, pain during activity, and amount of pain evoked during activity). Plaster casts appeared to be more effective in reducing elbow pain in tennis players compared to acupuncture and physiotherapy. Previous research supports these findings. Tonks et al. (47) observed significant pain reduction and



functional improvement in groups receiving topical corticosteroid injections, with these benefits persisting at the 3-month and 6-month follow-up assessments. However, Shakoori et al. (48) reported that only pain intensity, and not other aspects of pain, significantly decreased in the injection group compared to the phonophoresis group during their study period. Several studies support the use of corticosteroids for pain relief in certain conditions.

Newcomer et al. (49) found no significant difference in clinical outcomes between corticosteroid and placebo groups, but the corticosteroid group did experience a significant reduction in pain severity. Arti et al. (50) reported that the corticosteroid-treated group showed the highest increase in grip strength, the greatest pain relief, and the largest decrease in pain scores compared to other treatment groups at the first follow-up. However, it's important to note that the phrasing used in the study by Arti et al. regarding grip strength scoring might require clarification (81% grip strength (respectively), Score 1) and 83% of the healthy side (score 1), score against 4 and final score of 6). Jansen et al. (51) found that 30% of patients achieved permanent and complete pain relief after receiving a steroid injection and taking anti-inflammatory drugs. This study suggests that corticosteroid injection and casting may be more effective in promoting recovery for patients with tennis elbow compared to acupuncture and physiotherapy.

This study found that corticosteroid injection and casting had a greater effect on hand movement compared to acupuncture and physiotherapy. Patients in the corticosteroid and casting group achieved greater range of motion in terms of supination, pronation, extension, and flexion than those in the acupuncture and physiotherapy group. Jansen et al. (51) support these findings. Their study showed that corticosteroid injection and casting in patients with tennis elbow reduced wrist extensor activity, lessened pain,

and improved grip strength in both flexion and extension.

Several studies support the use of conservative physiotherapy for tennis elbow. Bisset et al. (32) and Emanet et al. (52) found that these approaches, including laser therapy, shockwave therapy, and exercise, offer longer-lasting therapeutic effects and fewer side effects compared to corticosteroid injections. However, it's important to note some limitations in the study by Arti et al. (50). While they report an initial average grip strength of 78% (score 3) and 47% (score 1) in flexion and extension of the affected elbow, respectively, they also state that scores decreased from treatment in all groups. This suggests the scoring system used might require further explanation.

Several studies have investigated the effectiveness of topical corticosteroid injections for tennis elbow. Assendelf et al. (24) reported that these injections were relatively safe and provided short-term benefits (lasting 2 to 6 weeks). Other research supports these findings, indicating that topical corticosteroids can reduce pain, improve overall recovery, and even enhance punching power in patients compared to placebo or other supportive treatments (23, 53). However, studies like the one by Kim et al. (25) suggest a potential limitation: while these injections may reduce pain in the short term (around 2 weeks) compared to physiotherapy, they might not have a lasting impact on overall patient outcomes by the 6-week mark.

Multiple studies suggest that a combination of oral NSAIDs and physiotherapy offers more advantages in the medium-term (beyond 6 weeks) and long-term (beyond 6 months) compared to corticosteroid injections (27, 53, 54). While corticosteroid injections boast a higher success rate in the short term (92% vs. 47% for physiotherapy), they are more likely to lead to recurrences in the long run (53). Additionally, nearly all patients treated with steroid injections experience a

return of severe symptoms at some point (8).

Smidt et al. (53) conducted a study involving 185 patients with lateral epicondylitis (tennis elbow). The study compared three treatment options: topical steroid injections, physiotherapy, and a no-intervention control group. The researchers concluded that physiotherapy offered the best long-term treatment strategy, while topical steroid injections provided a more pronounced effect in the short term.

Uygur et al. (55) investigated the use of dry needling compared to corticosteroids for treating lateral epicondylitis (tennis elbow). Their findings showed that both treatments were effective. However, the study suggests that dry needling may lead to significantly greater improvements in PRTEE scores at the 6-month mark compared to corticosteroid therapy. While corticosteroids offer short-term pain relief, repeated injections should be avoided. This is because they may not only damage tendons in the long term but also potentially reduce the success rate of surgery.

Several studies have investigated the effectiveness of acupuncture for treating tennis elbow. For example, Affaitati et al. (56) compared acupuncture to other treatments. In their study, acupuncture needles were inserted into trigger points located in the extensor muscles of the inflamed area (lateral epicondylitis). A placebo needle was also used as a control to compare the effects of the actual acupuncture treatment.

Langevin et al. (57) investigated the use of dry acupuncture for treating pain in specific shoulder muscles: the muscles under the spine, the upper trapezius, latissimus dorsi (large round muscle), and anterior deltoid. Their study found that dry acupuncture significantly reduced pain in these muscles, with reductions of 77%, 58%, 49%, and 38%, respectively. Dry acupuncture is thought to work by stimulating the release of endorphins, the body's natural pain relievers (58). In a study, Rothschild et al. found that dry acupuncture in tight muscle bands relieved chest

pain in the spine, neck, shoulders, and chest, although further studies are needed to confirm these findings (59). Dry acupuncture reduces muscle pain by affecting the activity of trigger points. It seems that with effective treatment of myofascial trigger points in the extensor muscles, the strain on the extensor muscles is reduced and the patient's pain is consequently reduced. The tight bands in the extensor muscles lead to chronic strain on the tendon joints of the extensors, which ultimately worsens the patient's symptoms.

The only limitation of the present study is the lack of complete cooperation between patients and incomplete hospital records, which according to the evaluation of the study over a long period has attempted to solve the problem. Because the present study was performed on patients in Jiroft with a small sample size and studies in this field are few, further studies with a larger sample size and a longer follow-up period are needed to confirm the effects of these methods and determine if they should be chosen as the preferred treatment method.

### Conclusion

Based on the findings of this study and previous research, corticosteroid injections with casting may be a more effective treatment for tennis elbow compared to acupuncture and physiotherapy. Additionally, corticosteroid injections appear to be effective in reducing pain associated with tennis elbow.

### Acknowledgments

This research originated from the dissertation of a Doctor of Medicine student at Jiroft University of Medical Sciences. The study was registered with the ethics committee under code IR.JMU.RE.1400.031 and received financial support from the university. The authors gratefully acknowledge the Vice-Chancellor for Research, the esteemed professors, advisors, and

all those who contributed to this research.

### **Conflict of Interest**

The authors declare that there is no conflict of interest in this study. We certify that this submission is a original work, and is not currently under consideration for publication in any other journal.

### **Funding**

We did not receive any funding for the present study.

### **Ethical Considerations**

This study was approved by the Research Ethics Committee of Jiroft University of Medical Sciences; Jiroft , Iran.

### **Code of Ethics**

It was registered under the ethical code IR.JMU.RE.1400.031 and financially supported by the university.

### **Authors' Contributions**

Conceived and designed the study: SH, RF and RR. Performed the experiments: MG, AA, SH and SD. Analyzed the data: SD and AJ. Contributed reagents/materials/analysis tools: FAG, SD and AJ. Drafted and revised the manuscript: SD and AJ.

### **Data Availability Statement**

Data are available on request from the authors.

### **List of Abbreviations**

VAS: Visual Analogue Scale

ANOVA: Analysis of variance

### **Declarations**

Ethical approval and informed consent: This study was conducted in accordance with the ethical standards of the institutional and/or national research committee and with the Declaration of Helsinki and its later amendments or comparable ethical standards. No animal

studies were conducted by the authors for this research.

### **References**

1. Matache BA, Berdusco R, Momoli F, Lapner PL, Pollock J. A randomized, double-blind sham-controlled trial on the efficacy of arthroscopic tennis elbow release for the management of chronic lateral epicondylitis. *BMC musculoskeletal disorders*. 2016;17(1):1-8.
2. Nirschl RP, Ashman ES. Tennis elbow tendinosis (epicondylitis). *Instructional course lectures*. 2004;53:587-98.
3. Coombes BK, Bisset L, Vicenzino B. Thermal hyperalgesia distinguishes those with severe pain and disability in unilateral lateral epicondylalgia. *The Clinical journal of pain*. 2012;28 (7):595-601.
4. Viswas R, Ramachandran R, Korde Anantkumar P. Comparison of the effectiveness of supervised exercise program and Cyriax physiotherapy in patients with tennis elbow (lateral epicondylitis): a randomized clinical trial. *The scientific world journal*. 2012;2012.
5. Kazemi M, Azma K, Tavana B, Moghaddam FR, Panahi A. Autologous blood versus corticosteroid local injection in the short-term treatment of lateral elbow tendinopathy: a randomized clinical trial of efficacy. *American journal of physical medicine & rehabilitation*. 2010;89(8):660-7.
6. Cutts S, Gangoo S, Modi N, Pasapula C. Tennis elbow: A clinical review article. *Journal of orthopaedics*. 2020;17:203-7.
7. Zeisig E, Öhberg L, Alfredson H. Extensor origin vascularity related to pain in patients with tennis elbow. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2006;14(7):659-63.
8. Santiago AO, Rios-Russo JL, Baerga L, Micheo W. Evidenced-based management of tennis elbow. *Current Physical Medicine and Rehabilitation Reports*. 2021:1-9.
9. Edwards SG, Calandruccio JH. Autologous blood injections for refractory lateral epicondylitis. *The Journal of hand surgery*. 2003;28(2):272-8.
10. Suresh S, Ali KE, Jones H, Connell D. Medial epicondylitis: is ultrasound guided autologous blood injection an effective treatment? *British journal of sports medicine*. 2006;40(11):935-9.
11. Xu B, Goldman H. Steroid injection in lateral epicondylar pain. *Australian Journal of General Practice*. 2008;37(11):925.
12. Kemp JA, Olson MA, Tao MA, Burcal CJ. Platelet-rich plasma versus corticosteroid injection for the treatment of lateral epicondylitis: a systematic review of systematic reviews. *International Journal of Sports Physical Therapy*. 2021;16(3):597.



13. Bisset L, Coombes B. Tennis elbow. *BMJ clinical evidence*. 2011;2011.
14. Hadeed MM, MacDonell JR, Dempsey IJ, Moore CC, Browne JA. Chronic Nocardia cyriacigeorgica Periprosthetic Knee Infection Successfully Treated with a Two-Stage Revision: A Case Report. *JBJS case connector*. 2017;7(4):e74.
15. Kalawadia JV, Kalainov DM. Tennis Elbow: Complications of Surgical Treatment and Salvage Procedures for Failed Surgery. *Tennis Elbow: Springer*; 2015. p. 153-67.
16. Brummel J, Baker III CL, Hopkins R, Baker Jr CL. Epicondylitis: lateral. *Sports medicine and arthroscopy review*. 2014;22(3):e1-e6.
17. Asghari E, Zarifian A, Shariyate MJ, Kachooei AR. Perceived Pain Severity and Disability After the Recurrence of Tennis Elbow Following a Local Corticosteroid Injection. *Archives of Bone and Joint Surgery*. 2022;10(9):760.
18. McMillan AM, Landorf KB, Gilheany MF, Bird AR, Morrow AD, Menz HB. Ultrasound guided corticosteroid injection for plantar fasciitis: randomised controlled trial. *BMJ*. 2012;344.
19. Mishra A, Pavelko T. Treatment of chronic elbow tendinosis with buffered platelet-rich plasma. *The American journal of sports medicine*. 2006;34(11):1774-8.
20. Barrett BSL, Erredge SE. Feature: growth factors for chronic plantar fasciitis. *Podiatry Today-ISSN*. 2004;17:1045-7860.
21. James SL, Ali K, Pocock C, Robertson C, Walter J, Bell J, et al. Ultrasound guided dry needling and autologous blood injection for patellar tendinosis. *British journal of sports medicine*. 2007;41(8):518-21.
22. Klein MB, Yalamanchi N, Pham H, Longaker MT, Chan J. Flexor tendon healing in vitro: effects of TGF- $\beta$  on tendon cell collagen production. *The Journal of hand surgery*. 2002;27(4):615-20.
23. Assendelft W, Green S, Buchbinder R, Struijs P, Smidt N. Tennis elbow Clin Evid. 2004; 1633-44. lateral epicondylitis. *Phy Ther Sci*. 2011;23:624-29.
24. Nambi G, Alghadier M, Verma A, Aldhafian OR, Alshahrani NN, Saleh AK, et al. Clinical and radiological effects of Corticosteroid injection combined with deep transverse friction massage and Mill's manipulation in lateral epicondylalgia—A prospective, randomized, single-blinded, sham controlled trial. *Plos one*. 2023;18(2):e0281206.
25. Struijs PA, Smidt N, Arola H, Van Dijk C, Buchbinder R, Assendelft WJ. Orthotic devices for the treatment of tennis elbow. *Cochrane Database of Systematic Reviews*. 2002(1).
26. Akkurt S, Yilmaz A, Saka T. A comparison of extracorporeal shock wave therapy, physiotherapy, and local steroid injection in treatment of lateral epicondylitis. *Turk J Phys Med Rehab*. 2016;1:37-44.
27. Crider K, Williams J, Qi YP, et al. Folic acid supplementation and malaria susceptibility and severity among people taking antifolate antimalarial drugs in endemic areas. *The Cochrane Database of Systematic Reviews*. 2022;2(2022). DOI: 10.1002/14651858.cd014217. PMID: 36321557; PMCID: PMC8805585.
28. Clarke H, Kirkham KR, Orser BA, Katznelson R, Mitsakakis N, Ko R. Gabapentin reduces preoperative anxiety and pain catastrophizing in highly anxious patients prior to major surgery: a blinded randomized placebo-controlled trial. *Canadian Journal of Anesthesia/Journal canadien d'anesthésie*. 2013;60(5):432-43.
29. Kongmalai P, Chanlalit C. Demographic causes of chronic lateral elbow pain along arthroscopic criteria. *J Med Assoc Thai*. 2016;99(8):S79-S83.
30. Shinde SB, Varadharajulu G. Effect of therapeutic exercise programme in adults with early rheumatoid arthritis. *Indian J Physiother Occup Ther*. 2017;11:76-80.
31. Green S, Buchbinder R, Barnsley L. Acupuncture for lateral elbow pain. *The Cochrane Database of Systematic Reviews*. 2002 (1):CD003527. DOI: 10.1002/14651858.cd003527. PMID: 11869671; PMCID: PMC8717066.
32. Bisset L, Paungmali A, Vicenzino B, Beller E. A systematic review and meta-analysis of clinical trials on physical interventions for lateral epicondylalgia. *British journal of sports medicine*. 2005;39(7):411-22.
33. Loew LM, Brosseau L, Tugwell P. Deep transverse friction massage for treating lateral elbow or lateral knee tendinitis. *The Cochrane Database of Systematic Reviews*. 2014 (11):CD003528. DOI: 10.1002/14651858.cd003528.pub2. PMID: 25380079; PMCID: PMC7154576.
34. Dabholkar AS, Kalbande VM, Yardi S. Neural tissue mobilisation using ULTT2b and radial head mobilisation v/s exercise programme in lateral epicondylitis. *Indian Journal of Physiotherapy and Occupational Therapy*. 2013;7(4):247.
35. Meunier M. Lateral epicondylitis/extensor tendon injury. *Clinics in sports medicine*. 2020;39(3):657-60.
36. Hong QN, Durand M-J, Loisel P. Treatment of lateral epicondylitis: where is the evidence? *Joint Bone Spine*. 2004;71(5):369-73.
37. Murtezani A, Ibraimi Z, Villasolli TO, Sllamniku S, Krasniqi S, Vokrii L. Exercise and therapeutic ultrasound compared with corticosteroid injection for chronic lateral epicondylitis: a randomized controlled trial. *Ortop Traumatol Rehabil*. 2015;17(4):351-7.
38. Silverstein B, Viikari-Juntura E, Kalat J. Use of a prevention index to identify industries at high risk for work-related musculoskeletal disorders of the neck, back, and upper extremity in Washington state, 1990–1998. *American journal of industrial medicine*. 2002;41(3):149-69.

39. Di Filippo, L, Vincenzi, S, Pennella, D, Maselli, F. Treatment, Diagnostic Criteria and Variability of Terminology for Lateral Elbow Pain: Findings from an Overview of Systematic Reviews. *Healthcare*. 2022, 10, 1095. <https://doi.org/10.3390/healthcare10061095>
40. Jiménez I, Muratore-Moreno G, Marcos-García A, Medina J. *Revista Española de Cirugía Ortopédica y Traumatología*. 2016.
41. Dimberg L. The prevalence and causation of tennis elbow (lateral humeral epicondylitis) in a population of workers in an engineering industry. *Ergonomics*. 1987;30(3):573-9.
42. Kim YJ, Wood SM, Yoon AP, Howard JC, Yang LY, Chung KC. Efficacy of nonoperative treatments for lateral epicondylitis: a systematic review and meta-analysis. *Plastic and Reconstructive Surgery*. 2021;147(1):112-25.
43. Mandal P, Chattopadhyay P. Treatment of Refractory Lateral Epicondylitis of Humerus with Local Autologous Plasma Injection. *International Journal of scientific study*. 2016;4(9):83-7.
44. Kivi P. The etiology and conservative treatment of humeral epicondylitis. *Scandinavian journal of rehabilitation medicine*. 1983;15(1):37-41.
45. Hutson MA. *Work-related upper limb disorders: Recognition and management*: Butterworth-Heinemann; 1997.
46. Rayan F, Rao VS, Purushothamdas S, Mukundan C, Shafqat SO. Common extensor origin release in recalcitrant lateral epicondylitis: role justified? *Journal of orthopaedic surgery and research*. 2010;5(1):1-3.
47. Tonks J, Pai S, Murali S. Steroid injection therapy is the best conservative treatment for lateral epicondylitis: a prospective randomised controlled trial. *International journal of clinical practice*. 2007;61(2):240-6.
48. Shakouri SK, Eftekhari-Sadat B, Johari F, Ghojzadeh M, Hemmati S, Ghaffari G. To Compare the Effect of Ultrasound Guided Local Injection of Dexamethasone with Dexamethasone Phonophoresis for Treatment of Lateral Epicondylitis (Tennis Elbow). *Journal of Arak University of Medical Sciences*. 2015;18(9):57-65.
49. Newcomer KL, Laskowski ER, Idank DM, McLean TJ, Egan KS. Corticosteroid injection in early treatment of lateral epicondylitis. *Clinical journal of Sport medicine*. 2001;11(4):214-22.
50. Arti H, Abrishamkar S, Rafieian M. Comparative Results of Non-surgical Long Term Treatments of Tennis Elbow Disease. *Journal of Advances in Medical and Biomedical Research*. 2005;13(50):1-7.
51. Jansen CWS, Olson SL, Hasson SM. The effect of use of a wrist orthosis during functional activities on surface electromyography of the wrist extensors in normal subjects. *Journal of Hand Therapy*. 1997;10(4):283-9.
52. Emanet SK, Altan LI, Yurtkuran M. Investigation of the effect of GaAs laser therapy on lateral epicondylitis. *Photomedicine and laser surgery*. 2010;28(3):397-403.
53. Smidt N, Assendelft WJ, van der Windt DA, Hay EM, Buchbinder R, Bouter LM. Corticosteroid injections for lateral epicondylitis: a systematic review. *Pain*. 2002;96(1-2):23-40.
54. Katz J, Melzack R. Measurement of pain. *Surgical Clinics of North America*. 1999;79(2):231-52.
55. Uygur E, Aktaş B, Yilmazoglu EG. The use of dry needling vs. corticosteroid injection to treat lateral epicondylitis: a prospective, randomized, controlled study. *Journal of Shoulder and Elbow Surgery*. 2021;30(1):134-9.
56. Affaitati G, Costantini R, Fabrizio A, Lapenna D, Tafuri E, Giamberardino MA. Effects of treatment of peripheral pain generators in fibromyalgia patients. *European journal of pain*. 2011;15(1):61-9.
57. Langevin HM, Bouffard NA, Badger GJ, Churchill DL, Howe AK. Subcutaneous tissue fibroblast cytoskeletal remodeling induced by acupuncture: Evidence for a mechanotransduction-based mechanism. *Journal of cellular physiology*. 2006;207(3):767-74.
58. Niddam DM, Chan R-C, Lee S-H, Yeh T-C, Hsieh J-C. Central representation of hyperalgesia from myofascial trigger point. *Neuroimage*. 2008;39(3):1299-306.
59. Rothschild B. Mechanical solution for a mechanical problem: Tennis elbow. *World journal of orthopedics*. 2013;4(3):103.