



## Original Article

## In Vitro Effects of Propolis Mouthwash and Nystatin on *Candida Albicans*: A Comparative Study

Alemrajabi Mohammad Sadegh<sup>1</sup>, Neshati Ammar<sup>2</sup>, Moosavi Vahid<sup>1</sup>, Sadrzadeh Afshar Maryam Sadat<sup>\*</sup>

1. Oral &amp; Maxillofacial Medicine Department, Faculty of Dentistry, Aja University of Medical Sciences, Tehran, Iran

2. Prosthodontics Department, Faculty of Dentistry, Aja University of Medical Sciences, Tehran, Iran

Received: 22 May 2021 Accepted: 30 Jun 2021

### Abstract

**Background & Objective:** Oral candidiasis is the most common fungal infection of the oral cavity, mainly caused by the overgrowth of *Candida albicans*. One of the topical medications for candidiasis is nystatin, a polyene antifungal agent. Nevertheless, increased resistance to this drug because of overprescription has caused recurrent oral candidiasis. There is a lot of interest today in the use of natural products and compounds due to the side effects of synthetic products. Propolis is a natural resin substance produced by bees through the combination of wax and saliva with resins collected from plants, which has reportedly antibacterial, antifungal and antitumor properties. The purpose of this study was to compare the effects of propolis mouthwash and nystatin on *C. albicans*.

**Materials & Methods:** The present in vitro study, in which the effects of antifungal agents were investigated on the standard strain of *C. albicans*, was conducted within two groups of 15, nystatin and propolis, using the Kirby-Bauer disc diffusion susceptibility test after preparation and sterilization. The diameter of the zone of inhibition (ZOI) was measured in this assay using the disc diffusion method onto Sabouraud dextrose agar (SDA) medium.

**Results:** Statistical analysis of data by SPSS software using t-test statistics to evaluate the antifungal effect of the interventions showed a better antifungal effect of nystatin compared to propolis mouthwash against *C. albicans* ( $P < 0.001$ ).

**Conclusion:** The findings of this study demonstrated that the local propolis mouthwash was less potent than nystatin in inhibiting the growth of *C. albicans*, probably due to differences in the concentration and geographical region of collected propolis. Further research on species isolated from oral biofilm is needed to achieve complementary outcomes.

**Keywords:** propolis mouthwash, nystatin, *Candida albicans*

### Introduction

*Candida albicans* is a fungus as a part of the normal microflora of the oral cavity in 20-50% of the healthy population, which is present in all surfaces of the oral mucosa. The presence of local or systemic predisposing factors causes the pathogenicity of this fungus (1).

**\*Corresponding Author: Sadrzadeh Afshar Maryam Sadat**, Oral & Maxillofacial Medicine Department, Faculty of Dentistry, Aja University of Medical Sciences, Tehran, Iran

Email: [m\\_sadrzade@alumnus.tums.ac.ir](mailto:m_sadrzade@alumnus.tums.ac.ir)<https://orcid.org/0000-0001-8277-3805>

*C. albicans* is known as the most important organism in the pathogenesis of oral candidiasis (2-4). Antifungals can be grouped into several classes depending on their mechanism of action, each of which affects different parts of the fungus, including cell membrane, genetics and cell divisions. Nystatin is an antibiotic of the polyene family that binds to the sterols in the fungal cell membrane, altering the membrane permeability and causing the release

Alemrajabi Mohammad Sadegh: <https://orcid.org/0000-0001-9221-0686>Neshati Ammar: <https://orcid.org/0000-0003-4732-2157>Moosavi Vahid: <https://orcid.org/0000-0003-4736-9246>



of essential intracellular contents of the fungus. Nystatin is given as a topical ointment or oral form in the management of mucosal candidiasis, such as infections of the oral cavity and gastrointestinal tract, and the treatment of alopecia areata. At present, the main prescription of nystatin is in the treatment of superficial and cutaneous infections, so that the nystatin ointment is applied directly to the infected site (5). Appropriate clinical responses to treatment are affected by several factors, including the individual's immune system, the susceptibility of the pathogen to the drug, how the drug penetrates and distributes, and the acceptance of the drug by the host, and the presence or absence of an infectious agent. Opportunistic microorganisms will grow and multiply if the host conditions change and the immune systems weaken. Conditions are created for the overgrowth of opportunistic agents under the influence of factors such as the administration of broad-spectrum antibiotics, the use of invasive methods of diagnosis and treatment, immune system defects (such as AIDS and pneumonia) and the use of immunosuppressive drugs in diseases like cancers. Another parameter associated with opportunistic infections is the presence of polymorphism in some of these agents (such as the genus *Candida*), which are resistant to inhibitory factors (6). Fungal infections caused by *C. albicans*, especially non-*albicans* *Candida* (NAC) species, have increased in recent years. Most NAC species have shown greater resistance to antifungals and elevated mortality rates in patients with candidiasis (7). Drug resistance, or antimicrobial resistance, occurs when agents such as bacteria, viruses, parasites, and fungi change in such a way those drugs previously used to treat infections become ineffective. When microorganisms become resistant to most antimicrobials, they can cause death, transmission to others, and high costs to individuals and society (8).

Polyenes target ergosterols, the essential component of fungal cell membranes. The number of yeast species resistant to some polyenes, such as amphotericin B and nystatin,

has increased recently (9, 10).

Various antifungals such as nystatin, clotrimazole, amphotericin B are commonly prescribed as the first line of treatment. However, these treatments are associated with side effects such as bitter taste, allergic reactions, adrenal insufficiency, liver necrosis, drug interactions and resistance, which potentially limit their antifungal benefits (11-14). The increasing incidence of clinical resistance to antifungal therapy and the lack of response to it in recent years emphasize the need to introduce new treatments and additional prevention and treatment candidates (15). Accordingly, there is a need to increase general knowledge of natural products; however, the use of these natural products is limited due to scattered research (16-18). Hence, today's global attitude is to reduce the administration of synthetic products and the exploitation of traditional medicine (19). Propolis is a naturally occurring resin substance produced by bees through the combination of wax and saliva with resins collected from plants (20). The propolis has been shown to have antibacterial, antifungal and antitumor properties. The composition of propolis varies in different regions due to differences in plants of various regions, which causes diverse properties of this substance (21, 22). Propolis extract contains a wide range of components such as flavonoids and phenolic acids. The flavonoids in propolis, mainly Pinocembrin, are responsible for the inhibition of *Candida* (23). Due to the prevalence of fungal diseases, side effects of nystatin administration and differences in propolis compounds (thus its antifungal properties) in different regions, the present in vitro study aimed to compare the antifungal effects of local propolis mouthwash and nystatin on *C. albicans*.

### **Materials & Methods**

The current study was conducted on *C. albicans* prepared from the mycology center of Tehran University of Medical Sciences, Iran (Table 1) (Fig 1-A). Propolis-based mouthwash was native to the city of Mashhad, Iran (Soren Tech Toos, Mashhad, Iran) (Fig1-B). The effective dose of mouthwash was 4 mg/mL. The

nystatin discs were 100 IU/mL (Padtan Teb, Tehran, Iran) (Table 2) (Fig 1-C). The culture medium was Sabouraud dextrose agar (SDA), prepared according to the instructions on the medium and then autoclaved (at a temperature of 91°C and a pressure of 121 for 15 minutes) (Fig 1-D). After cooling, the sterile medium was distributed in fifteen 10 cm plates and stored in the refrigerator. The therapeutic concentration (without dilution) of propolis mouthwash was then poured into an empty sterile plate. Next, 15 sterile blank paper discs with a diameter of 4.6 mm were taken by sterile forceps and immersed in the mouthwash-containing plate to be completely wet. After draining the excess solution, the plates were placed in an oven at a temperature of 40°C for about 20 minutes to evaporate the excess mouthwash solutions to dry the discs (24) (Fig 1-E). After cooling the medium, the fungal single colony was spread onto the medium by sterile swab under sterile conditions. The dried discs were carefully seeded onto the medium inside the plate (containing fungal culture) via sterile forceps at a distance


of 10 to 15 mm from the edge of the plate and gently pressed on the agar surface thus that the whole disc was in contact with the agar. The nystatin discs with a diameter of 4.6 mm were also planted by sterile forceps carefully onto the medium inside the plate (containing fungal culture) at a distance of 10 to 15 mm from the edge of the plate and gently pressed on the agar surface until the whole disc was in contact with the agar (Fig 1-F). Incubation was performed at 37°C for 24 and 48 hours. Finally, the diameter of the zone of inhibition (ZOI) was measured in millimeters by a caliper (with an accuracy of 0.1 mm) and recorded in a pre-prepared form. The diameter of the ZOI appearing around the nystatin and propolis mouthwash discs against *C. albicans* was carefully measured under the desk lamp and compared by analysis of variance (ANOVA) test. The SDA medium was applied to isolate a single colony and the disc diffusion method to determine the susceptibility of fungus to mouthwash and nystatin (25). It is noteworthy that all the above-mentioned steps were carried out aseptically under the hood and in the vicinity of the flame (Fig 1-G).

**Table 1.** Studied microbial sample properties

Microbial sample name	Common culture medium	Incubation suitable temperature	Incubation time
<i>Candida Albicans</i>	Sabouraud dextrose agar	37° C	24 & 48 hours

**Table2.** Studied antimicrobial agents

Antimicrobial agent	Manufacturer	Country-city of manufacture	Product Code
Propolis Mouthwash	Soren Tech Toos	Iran, Mashhad	557310
Nystatine disc	Padtan Teb	Iran, Tehran	132464



**Figure 1.** A) *Candida Albicans*. B) Propolis mouthwash. C) Nystatine discs. D) Prepared culture medium. E) Mouthwash impregnated discs. F) culture mediums containing Nystatine and Propolis. G) working under the hood and in the vicinity of the flame. H) ZOI in culture mediums

**Results**

In the statistical analysis of the effect of propolis mouthwash and nystatin on *C. albicans*, two different effects were evaluated: 1- the effect of propolis mouthwash on ZOI, and 2- the effect of nystatin on ZOI (Table 3) and (Fig 1-H). The method of data collection in this study was a checklist (Table 4). In data analysis, first, the normal distribution of data was investigated by Chi-square test and the data were expressed as SEM ± Mean and analyzed by unpaired t-test (P<0.05).

According to the relevant variables and the mean ZOI diameter (SEM ± Mean), the results showed that the nystatin was more effective than propolis mouthwash on *C. albicans* and the propolis mouthwash had little effect on *C. albicans* (P<0.001). (Chart 1)


The diameter of the zone of inhibition in the checklist above was measured in millimeters using the caliper.

**Table 3.** Descriptive Statistics of the effect of the agents on C.albicans

Anti-fungal Agent	Number	Mean	Std.Error Mean	P_value
Nystatine	15	10.53	0.375	0.000
Propolis	15	0.53	0.140	

**Table 4.** check list of ZOI in millimeters in each of agents

Fungus	<i>C. Albicans</i>					P_value
Anti-Fungal agent						
Nystatine	11	12	12.5	8.5	12	<0.01
	12	10	10.5	10.5	11	
	9	8	10	11	10	
Propolis	2	1	1	0	0.5	
	0.5	1	0.5	0	1	
	0.5	0	0.5	0	0.5	



**Chart 1.** ZOI in millimeter in antifungal agents. (Nytatin: 10.53 mm and Propolis 0.53)



## Discussion

Oral candidiasis is the most common fungal infection of the oral cavity, so it is clear that studies are needed to find a solution to overcome this disease. Following the spread of AIDS and the high prevalence of various cancers, the incidence rate of opportunistic oral infections, such as fungal infections, has increased due to a weakened immune system, the nature of the disease itself or immunosuppressants used in treatment. Finding the right strategy to solve such problems is also necessary in these special cases. Prescribing antiseptics and disinfectants, including mouthwashes, is one of these approaches (26). Various antifungals such as nystatin, clotrimazole, amphotericin B and chlorhexidine are commonly administered as the first line of treatment for candidiasis. However, these treatments are associated with side effects such as bitter taste, allergic reactions, adrenal insufficiency, liver necrosis, drug interactions, and resistance, which potentially limit their antifungal benefits. Despite recent advances and new technologies in the fabrication of health products, there is a great deal of interest in using natural products and compounds due to the side effects of synthetic products, and recent research has focused on investigating the properties and feasibility of their widespread application (16 -18). Propolis is a natural plant-derived resin produced by bees from the flowers, pollen, branches and leaves of plants and used to repair the hive wall and protect the colony against disease (20). Propolis has long been prescribed to heal oral ulcers (27) and has been shown to have antibacterial, antifungal, antiviral, antioxidant, antitumor, and anti-inflammatory properties (28). Propolis extract contains a wide range of components such as flavonoids and phenolic acids. The flavonoids in propolis, mainly pinocembrin, account for the inhibition of *Candida*. These mouthwashes are better accepted by Iranian patients due to their availability in the country, low cost, acceptable taste and smell, easy use, and non-chemical nature. The fungus studied in this study is part of the microflora of the oral cavity and is involved in the occurrence

of oral diseases and is found in all surfaces of the oral mucosa. *C. albicans* is known as the most important organism in the pathogenesis of candidiasis. Therefore, the aim of this in vitro study was to compare the effect of propolis mouthwash and nystatin on *C. albicans* (23). In this study, the ZOI diameter was measured using disc diffusion method onto the SDA medium. This method is based on the drug diffusion in the medium in the vicinity of the organism. In this method, the paper discs impregnated with a specific concentration of the drug were seeded onto SDA medium on which the fungal sample had grown. The results were read after incubation at 37°C for 24 and 48 hours by measuring the ZOI diameter, which is radically formed around the disc, and the susceptibility of the yeasts was evaluated. The results of disc diffusion method and ZOI diameter in this study showed that the efficacy of nystatin was higher than propolis mouthwash on *C. albicans* and propolis mouthwash had little effect on this fungus. The results of this study also revealed that the nystatin had the highest ZOI diameter, or in other words *C. albicans* exhibited the highest sensitivity to the nystatin (Table 3), which can be attributed to the concentration of commercial propolis mouthwash ( $P < 0.001$ ). According to the data table, the antifungal effect of nystatin is about 20 times higher than that of propolis mouthwash, and commercial propolis mouthwash has a limited effect on *C. albicans*. Therefore, propolis mouthwash can act against any microorganism in different ways. The difference between the ZOI diameters for nystatin and propolis mouthwash in this method and those found in other studies is probably due to differences in *C. albicans* tolerance or the origin of propolis samples, as the propolis composition depends on the regional vegetation (23). According to the studies of Silva et al. (29), and Prospero et al. (22), the differences in plants of various regions lead to diverse compositions of propolis in different regions, which causes variable properties of the substance. The lower efficiency of propolis mouthwash than nystatin in disc diffusion method on *C. albicans* can be



due to the constant concentration of mouthwash (with the active ingredient of 4 mg/mL). Therefore, in addition to the cost, if other methods, such as minimum inhibitory concentration (MIC), were performed, fewer efficacies would be seen for this mouthwash compared to nystatin, again due to the presence of a constant concentration of active ingredient. Diba et al. reported a great mean MIC value (30) of alcoholic propolis extract on the growth of *Candida* and *Aspergillus* isolates (31). Ganjavi et al. found that the propolis extract with the MIC value of 90 and the minimum fungicidal concentration (MFC) values of 39 and 65 µg/mL had the highest antifungal activity compared to other extracts studied, respectively. In addition, nystatin and amphotericin B showed better effects on laboratory fungi than the impact of all extracts studied on *C. albicans* (32). Herrer et al. found that all propolis samples inhibited *Candida* species; however, there were significant differences in extract concentrations, indicating inhibition of *Candida* species (33).

Pereira et al. documented evidence of the clinical efficacy of an alcohol-free mouthwash containing 5.0% Brazilian green propolis in the control of plaque and gingivitis (34). Silici et al. concluded that the bee products, especially propolis and pollen, could help control some fluconazole-resistant fungal species (35). According to the results of these studies, the difference in concentrations of propolis extract is one of the effective factors in determining the efficacy and inhibition of microbial species and this extract exerts its effect well at higher concentrations. Based on the studies by Santiago et al. (36), Mohan et al. (37), Acka et al. (38), the results suggest that the propolis mouthwash has stronger antifungal effects, and the reason goes back to the difference in various potential of mouthwashes and the difference in the formulation and properties of propolis studied, because the differences in the region, the season of propolis collection, its contamination with wax and bee species all lead to differences in the properties of propolis. On the other hand, the differences in the microbiological methods studied, including fungal species, cell differentiation phase, culture

and time conditions, duration of drug use and study design, are other causes of differences. The present study has advantages over previous studies, such as the use of local propolis; Minimal use of synthetic compounds and alcohol in mouthwash can reduce the effects of long-term use. Many articles have emphasized that geographical area affects the properties of propolis (38, 39).

### **Conclusion**

Considering the results of the study and less efficacy of propolis mouthwash over nystatin, it can be concluded that the treatment with propolis mouthwash can reduce oral diseases, especially fungal infections, although less than antifungals. Due to the stronger antifungal properties and far fewer side effects of herbal mouthwashes than chemical mouthwashes, they may be considered as potent antifungal agents, and since there is limited research on propolis mouthwash and its antimicrobial spectrum is unclear, the manufacturer is suggested increasing the concentration of the active ingredient to enhance the antifungal effect of this mouthwash. Further and extensive studies are needed before recommending the use of this mouthwash as an antifungal agent. According to the results of this study, it is important to note that the susceptibility of the pathogen to mouthwash should be measured in choosing an effective mouthwash to treat the infection, in addition to the type, concentration and antiseptic properties of the mouthwash, because the growth inhibitory effect depends on the type of microorganism, the type of mouthwash and the tested concentrations.

### **Statistically data analysis**

Data were statistically analyzed by SPSS version 2 software. First, the parametric or non-parametric nature of collected data was determined by the Kolmogorov-Smirnov test. Data were reported as Mean ± SEM and analyzed statistically by unpaired t-test student test and P<0.05 was considered as a significance level. Finally, the logistic regression test was performed for more accurate analysis. The



project was approved by the Ethics Committee of Aja University of Medical Sciences, Iran, under the code of ethics of IR.AJAUMS.REC.1399.023.

### **Acknowledgement**

The project was approved by the Ethics Committee of Aja University of Medical Sciences, Iran, under the code of ethics of IR.AJAUMS.REC.1399.023. The authors would like to thank the staff of Aja university of medical sciences who facilitated this study.

### **Conflict of Interest**

Authors declare no conflicts of interests associated with the publication of this article.

### **References**

- Mehdipour M, Hakemi Vala M, Sadrzadeh-Afshar MS, Gholizadeh N. In Vitro antifungal effect of cinnamon extract on candida species. *Caspian J Dent Res.* 2018; 7:49-53.
- Mehdipour M, Gholizadeh N, Sadrzadeh-Afshar M-S, Hematpoor N, Kalae P, Hakemi Vala M, et al. In Vitro Comparison of the Efficacy of Cumin Extract and Fluconazole against Candida Strains. *J Islam Dent Assoc Iran.* 2019; 31(2):98-107. DOI: 10.30699/jidai.31.2.98
- Bilhan H, Sulun T, Erkose G, Kurt H, Erturan Z, Kutay O, et al. The role of Candida albicans hyphae and Lactobacillus in denture-related stomatitis. *Clin oral investig.* 2009; 13(4):363.
- Webb B, Thomas C, Willcox M, Harty D, Knox K. Candida-associated denture stomatitis. Aetiology and management: A review: Part1. Factors influencing distribution of candida species in the oral cavity. *Aust dent J.* 1998;43(1):45-50.
- Barbeau J, Séguin J, Goulet JP, de Koninck L, Avon SL, Lalonde B, et al. Reassessing the presence of Candida albicans in denture-related stomatitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2003;95(1):51-9.
- White TC, Marr KA, Bowden RA. Clinical, cellular, and molecular factors that contribute to antifungal drug resistance. *Clin microbial rev.* 1998;11(2):382-402.
- Jahanshahi G, Yazdani M, Asadian F. An investigation on the relationship between oral candidal colony count and the duration of hemodialysis. *J of Dent Med.* 2003;16(3):46-51.
- Hill JA, Ammar R, Torti D, Nislow C, Cowen LE. Genetic and genomic architecture of the evolution of resistance to antifungal drug combinations. *PLoS Genet.* 2013;9(4):e1003390.
- Dick J, Merz W, Saral R. Incidence of polyene-resistant yeasts recovered from clinical specimens. *Antimicrob*

- Agents Chemother. 1980;18(1):158-63.
- Drutz D, Lehrer R. Development of amphotericin B-resistant *Candida tropicalis* in a patient with defective leukocyte function. *Am J med sci.* 1978;276(1):77-92.
- Chandra J, Mukherjee P, Leidich S, Faddoul F, Hoyer L, Douglas L, et al. Antifungal resistance of candidal biofilms formed on denture acrylic in vitro. *J dent res.* 2001;80(3):903-8.
- Ghapanchi J, Moattari A, Lavaee F, Shakib M. The antibacterial effect of four mouthwashes against *Streptococcus mutans* and *Escherichia coli*. *JPM.* 2015;65(350).
- Sanglard D, Coste A, Ferrari S. Antifungal drug resistance mechanisms in fungal pathogens from the perspective of transcriptional gene regulation. *FEMS yeast res.* 2009;9(7):1029-50.
- Yarborough A, Cooper L, Duqum I, Mendonça G, McGraw K, Stoner L. Evidence regarding the treatment of denture stomatitis. *J prosthodont.* 2016;25(4):288-301.
- Wise R, Hart T, Cars O. Antimicrobial resistance Is a major threat to public health (editorial). *BMJ.* 1998;317:609-10.
- Balappanavar AY, Sardana V, Singh M. Comparison of the effectiveness of 0.5% tea, 2% neem and 0.2% chlorhexidine mouthwashes on oral health: A randomized control trial. *Indian Dent Res.* 2013;24(1):26.
- Oprea TI, Tropsha A, Faulon J-L, Rintoul MD. Systems chemical biology. *Nat chem biol.* 2007;3(8):447-50.
- Coleman DC, O'Donnell MJ, Boyle M, Russell R. Microbial biofilm control within the dental clinic: reducing multiple risks. *J Infect Prev.* 2010;11(6):192-8.
- Khorakian F, Movahed T, Ghazvini K, Karbasi S, Tabrizi Nouri S, Bahramian L, et al. Evaluation of frequency of microbial contamination in clinical setting surface in Dental School of Mashhad University of Medical Sciences. *J Mashhad Dent Sch.* 2017;41(3):209-18.
- Tavafi, H, Sadrzadeh-Afshar, M-S, Niroomand, S. In vitro effectiveness of antimicrobial properties of propolis and chlorhexidine on oral pathogens: A comparative study. *Biosis.* 2020; 1(3):116-125. <https://doi.org/10.37819/biosis.001.03.0062>.
- Dettenkofer M, Wenzler S, Amthor S, Antes G, Motschall E, Daschner FD. Does disinfection of environmental surfaces influence nosocomial infection rates? A systematic review. *Am J of infect control.* 2004;32(2):84-9.
- Prospero E, Savini S, Annino I. Microbial aerosol contamination of dental healthcare workers' faces and other surfaces in dental practice. *Infect Control Hosp Epidemiol.* 2003;24(2):139-41.
- Metzner J, Schneidewind E, Friedrich E. Effect of propolis and pinocembrin on fungi. *Die Pharmazie.* 1977;32(11):730.
- Arbabi-Kalati F, Porzamani M. Comparison the antifungal effect of licorice and nystatin, invitro study. *J Dent Med.* 2013;26(1):71-4.
- Ansari Moghaddam S, Raeisi A, Mehrabani M,



- Ansarimoghaddam A. Evaluating the Effect of Propolis-containing Toothpaste on Dental Plaques of Dentistry Students of Dentistry School, Zahedan University of Medical Sciences, Zahedan, Iran. *Mashhad Dent Sch.* 2019;43(4):323-30.
26. Cabral RS, Alves CC, Batista HR, Sousa WC, Abrahão IS, Crotti AE, et al. Chemical composition of essential oils from different parts of *Protium heptaphyllum* (Aubl.) Marchand and their in vitro antibacterial activity. *Nat prod res.* 2019;1-6.
27. Dodwad V, Kukreja BJ. Propolis mouthwash: A new beginning. *J Indian Soc Periodontol.* 2011;15(2):121.
28. Ozan F, Sümer Z, Polat Z, Er K, Ozan U, Deger O. Effect of mouthrinse containing propolis on oral microorganisms and human gingival fibroblasts. *European Journal of Dentistry.* 2007;1(4):195-201.
29. Silva JC, Rodrigues S, Feás X, Estevinho LM. Antimicrobial activity, phenolic profile and role in the inflammation of propolis. *Food Chem Toxicol.* 2012;50(5):1790-5.
30. Santiago KB, Piana GM, Conti BJ, Cardoso EdO, Murbach Teles Andrade BF, Zanutto MR, et al. Microbiological control and antibacterial action of a propolis-containing mouthwash and control of dental plaque in humans. *Nat prod res.* 2018;32(12):1441-5.
31. Diba K, Mousavi B, Mahmoudi M, Hashemi J. In-vitro anti fungal activity of Propolis alcoholic extract on *Candida* spp. and *Aspergillus* spp. *Tehran Univ Med J.* 2010;68(2):80-6.
32. Gavanji S, Larki B. Comparative effect of propolis of honey bee and some herbal extracts on *Candida albicans*. *Chin J integ med.* 2017;23(3):201-7.
33. Herrera CL, Alvear M, Barrientos L, Montenegro G, Salazar LA. The antifungal effect of six commercial extracts of Chilean propolis on *Candida* spp. *Cie inv agr.* 2010;37(1):75-84.
34. Pereira EMR, da Silva JLDC, Silva FF, De Luca MP, Lorentz TCM, Santos VR. Clinical evidence of the efficacy of a mouthwash containing propolis for the control of plaque and gingivitis: a phase II study. *Evid based complement alternat med.* 2011 ;2011:750249. doi: 10.1155/2011/750249.
35. Silici S, Koç NA, Ayangil D, Çankaya S. Antifungal activities of propolis collected by different races of honeybees against yeasts isolated from patients with superficial mycoses. *J pharmacol sci.* 2005;99(1):39-44.
36. Santiago KB, Piana GM, Conti BJ, Cardoso EdO, Murbach Teles Andrade BF, Zanutto MR, et al. Microbiological control and antibacterial action of a propolis-containing mouthwash and control of dental plaque in humans. *Nat prod res.* 2018;32(12):1441-5.
37. Mohan PU, Uloopi K, Vinay C, Rao RC. In vivo comparison of cavity disinfection efficacy with APF gel, Propolis, Diode Laser, and 2% chlorhexidine in primary teeth. *Contemp clin dent.* 2016;7(1):45.
38. Akca AE, Akca G, Topçu FT, Macit E, Pıkdöken L, Özgen IŞ. The comparative evaluation of the antimicrobial effect of propolis with chlorhexidine against oral pathogens: An in vitro study. *BioMed res int.* 2016; Article ID 3627463.
39. Bazvand L, Aminozarbani MG, Farhad A, Noormohammadi H, Hasheminia SM, Mobasherizadeh S. Antibacterial effect of triantibiotic mixture, chlorhexidine gel, and two natural materials Propolis and Aloe vera against *Enterococcus faecalis*: An ex vivo study. *Dent res J.* 2014;11(4):469-74.