## Homepages http://jabsstums.acefr



# The Evaluation of Efflux Pump *Genes norA*, *norB* and *norC Related* to Fluoroquinolones Resistance in *Staphylococcus aureus* Strains Isolated from Blood Infection

Dalir Fatemeh¹<sup>10</sup>, Morovvati Abbas¹<sup>10</sup>, Javadi Ali²<sup>10</sup>, Afkhami Hamed³<sup>10</sup>, Khaledi Mansoor³<sup>10</sup>, Fathi Javad⁴<sup>10</sup>, Mohsenipour Zeinab⁵<sup>10</sup>, Dirbaziyan Ashkan¹<sup>10</sup>, Zargar Mohsen¹<sup>12</sup>

- 1. Department of Microbiology, Qom Branch, Islamic Azad University, Qom, Iran
- 2. Department of Medical Science, Qom Branch, Islamic Azad University, Qom, Iran
- 3. Department of Microbiology, School of Medicine, Shahed University, Tehran, Iran
- 4. Department of Bacteriology and Virology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran
- 5. Department of Microbiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

### **Article Info**

#### **Article Type:**

Original Article

#### **Article History:**

Received

04 Jan 2023

Received in revised form

11 Feb 2023

Accepted

03 Apr 2023

Published online

01 Mar 2023

#### **Publisher:**

Fasa University of Medical Sciences

### **Abstract**

**Background & Objective:** Recently, ciprofloxacin resistance in *Staphylococcus aureus* strains, due to efflux pumps, has become a significant challenge. Therefore, this study was performed to evaluate the frequency of *norA*, *norB*, and *norC* efflux pump genes and their roles in resistance to ciprofloxacin in clinical isolates of *S. aureus*.

**Materials & Methods:** A total of one hundred clinical blood samples were collected from patient in Qom hospitals and *S. aureus* isolates were identified by standard microbiological tests. Antimicrobial susceptibility patterns were determined by the disk diffusion method using CLSI guidelines. Subsequently, the presence of *norA*, *norB*, and *norC* efflux pump genes in ciprofloxacin isolates was detected using the PCR method.

**Results:** Among one hundred clinical samples, 36 *S. aureus* isolates were recovered and the results of antibiotic susceptibility tests showed that twenty of them were resistant to ciprofloxacin. 15 isolates were resistant to norfloxacin and one isolate was resistant to ofloxacin. Moreover, the *norA*, *norB*, and *norC* genes were found in 58%, 30%, and 41% of ciprofloxacin-resistant isolates, respectively.

**Conclusion:** Based on the results of this study, *norA*, *norB*, and *norC* efflux pumps may play a significant role in the development of resistance to ciprofloxacin in clinical isolates of *S. aureus*. Detecting these genes may prove useful in suggesting an effective treatment model for infections caused by *S. aureus*.

Keywords: Staphylococcus aureus, Blood infection, Bacterial drug-resistance, Efflux pump

Cite this article: Dalir F, Morovvati A, Javadi A, Afkhami H, Khaledi M, Fathi J, Mohsenipour Z, Dirbaziyan A, Zargar M. The Evaluation of Efflux Pump Genes *norA*, *norB* and *norC* Related to Fluoroquinolones Resistance in *Staphylococcus aureus* Strains Isolated from Blood Infection. JABS. 2023;13(1): 81-87. **DOI:** 10.18502/jabs.v13i1.12079

### **Introduction**

Antibiotic resistance has been recognized by the efflux pumps at *Staphylococcus aureus* for the past few decades, but its clinical relevance has only recently been discovered. Five families of these pumps have now been identified as Prokaryotes,

**Solution Solution Zargar Mohsen**, Department of Microbiology, Qom Branch, Islamic Azad University, Qom, Iran

Email: zmohsen2000@yahoo.com

leading to resistance to many antibiotics (1). Today, active efflux mechanisms are responsible for the resistance of bacteria to a variety of non-building antibiotics and toxic compounds, and it seems that reducing the permeability of bacteria is one of the primary reasons for explaining the resistance of bacteria to drugs (2). MFS carriers are probably the largest and most diverse pumps among all the families of efflux pumps



## Homepages http://jabs.fums.ac.fr



Zargar M, et al.

that have been found in all branches of biology (3). The QacA and norA pumps in Staphylococcus aureus belong to the family of MFS efflux pumps. Another family is the RND efflux pumps, which are single-component or multi-component transmission systems that contain not only an internal membrane transmitter, but also an external membrane channel and a compatible periplasmic protein (4). Efflux pumps are the most important cause of antibiotic resistance, and the most important of these pumps are the intrinsic efflux gene systems. In Staphylococcus aureus, norA, norB, norC, and tet38 are chromosomal genes that encode efflux pumps, and the high expression of multidrug resistance (MDR) genes can be resistant to Quinolones and other compounds or tetracycline (tet 38) (5). norA multifunctional efflux pump systems have been widely studied in Staphylococcus aureus. The chromosomal gene encoding the pump is norA, which was first identified and collected in 1986 at a hospital in Japan in resistant bacteria to fluoroquinolones (6). After initial studies of norA, evidence suggests that other efflux systems are present in the S. aureus chromosome. norB is another diffusion pump similar to norA, belonging to the MFS family and consisting of 463 amino acids with 12 membrane sections (7). norB is also resistant to some norA substrates and causes bacterial resistance to hydrophilic fluoroquinolones (such as norfloxacin and ciprofloxacin), pesticides, and ethidium bromide dye, as well as non-norA-dependent substrates, such as hydrophobic fluoroquinolones and moxifloxacin (8). Studies show that norB may be involved in the response of Staphylococcus aureus to acidic shock and cellular oxygen deprivation, conditions that increase norB gene expression. It has also been shown to play an important role in bacterial pathogenesis (9). In both cases, the increase in norB gene expression has been proven by increasing the resistance to the substrates of this pump. One of the important substrates of norB is moxifloxacin (10). Evidence suggests that norC efflux systems are present

in the S. aureus chromosome. norC is another

emission pump that is similar to *norA* in the MFS family. It consists of 462 amino acids with 12 membrane parts which is 61% similar to *norB*. This pump is less resistant to hydrophobic and hydrophilic fluoroquinolones such as ciprofloxacin, moxifloxacin and garinagsin, and rhodamine dye (11). One of the ways to deal with antimicrobial resistance caused by efflux pumps is to use new antimicrobial compounds or to use an accompanying compound called an antibiotic adjuvant (12).

The aim of this study was to evaluate efflux pump genes *norA*, *norB*, and *norC* related to fluoroquinolone resistance in *Staphylococcus* aureus strains isolated from blood infection.

### **Materials and Methods**

One hundred blood samples (13) were obtained from hospitalized patients in Qom hospitals. Samples were collected from patients hospitalized in different departments of Kamkar and Shahid Beheshti hospitals in 2019. Patients were selected for sampling who experienced continuous fever during the last 24 hours.

In order to identify the collected strains, gram staining, mannitol fermentation test, DNase, catalase, and coagulase were used. Then, sensitivity of isolates to Ciprofloxacin, Norfloxacin, and Ofloxacin antibiotics was determined using the Kirby-Bauer method. The results of the antibiogram were interpreted using the CLSI 2019 break-points (14).

DNA of confirmed isolates was purified by using a genomic DNA extraction kit (Sinaclon, Iran) as recommended by the manufacturer. 36 isolates were screened for the presence of *norA*, *norB*, and *norC* genes by PCR, using *norA* forward: GTGGTATGAGTGCTGGTATGG and reverse: GAAACTTCTGCCATAAATCCACC, norB forward: CAAACACTCGGATGCAAGAAAC and reverse: GACGCCAAATGCTCCACC, *norC* forward: TGGGTTGGAGATGGATTTTC and reverse: ACAATTAGCCCTGCAACGTC previously described primers (primers were designed in this study). For each reaction,

## Homepages https://jabs.fums.ac.fr



Efflux Pump Genes norA, norB and norC Related to Fluoroquinolones Resistance

a sample containing distilled water was used as a negative control and a confirmed isolate with the target gene (norA, norB, and norC) was used as a positive control.

PCR was down 33 cycles denaturation 95°C 4min, second denaturation 94°C 35 sec annealing 53°C 35 sec, extension 72°C 35 sec and final extension 72°C 5min. After purification, the OD of each of the positive samples was read and then the PCR product was sent to the Sinaclon Company to determine the sequence. The submitted sequences were analyzed with CLC Sequence Viewer software, then compared with standard gene-related sequences in the NCBI database and Finch Tv software (version 1.4.0).

Chi-square statistical test and SPSS-

26 software were used for statistical analysis of the data of this research.

### **Results**

Out of a total of one hundred samples collected, 36 strains of *Staphylococcus aureus* were isolated from 100 samples, and the morphological and biochemical features were the criteria for confirming the identity of the isolates. Antibiotic susceptibility of 36 isolates of *Staphylococcus aureus* was determined by the agar disk diffusion method to the desired antibiotics, which showed that out of 36 isolates of *Staphylococcus aureus* only 20 to ciprofloxacin disk (55%), 15 cases were resistant to norfloxacin (42%) and 1 case to ofloxacin (3%) (Chart 1).




Chart 1. the percentage of resistance to each antibiotic separately, noro; norfloxacin, cep: ciprofloxacin, ofl: ofloxacin

### Molecular test results

Out of one hundred samples collected, 36 samples had antibiotic resistance, and DNA extraction and PCR reaction were performed on them with specific primers after extracting the genomic DNA of bacteria. Electrophoresis of the samples in 2%

agarose gel in the initial PCR reaction showed a band of 160 bp *norA*, 260 bp *norB* and 130 bp *norC* which indicated the amplification of the genes (Figure 1). Out of 36 resistant samples, 58%, 30% and 41% of the samples showed a positive result based on primers and had an antibiotic resistance genes

## Homepages https://jabs.fums.ac.fr



Zargar M, et al.




Figure 1. nor A, nor B and nor C gel elechtrophoresis. SM: DNA ladder, NC: Negative control, PC: Positive control

Molecular analysis for the presence of *norA*, *norB*, and *norC* genes based on primers available for each gene showed that 58% of the isolates had *norA* gene. *norB* gene was present in 30% of the isolates and 41% of the

isolated strains had *norC* gene. Based on the obtained molecular results, 4% are positive for two genes and 11% of bacteria are positive for three genes. About 38% of the bacteria did not have any of the genes (Chart 2).




Chart 2. Frequency distribution of each gene

## Homepages https://jabs.fums.ac.fr



Efflux Pump Genes norA, norB and norC Related to Fluoroquinolones Resistance

### Results of the sequencing

The result of sequencing for strain 9 for *norA*, strain 14 for *norB* and 21 indicates a 100% similarity of these genes in *Staphylococcus aureus*.

### **Discussion**

The resistance of efflux pumps to fluoroquinolones, such as ciprofloxacin, norfloxacin, and Sparfloxacin, in clinical isolates of *Staphylococcus aureus* has been described in the last two decades (15-16).

In this study, resistance to norfloxacin was 42%, resistance to ciprofloxacin was 55% and resistance to ofloxacin was 3%. Molecular studies for the presence of *norA*, *norB* and *norC* genes were considered to have *norA* gene in 21 samples (58%) of isolated bacteria and in 11 samples (30%) of bacteria with primers of *norB* gene showed positive results. Also, in isolates for *norC* gene in 15 samples (41%) this gene was included. Based on the molecular results obtained in 4 bacteria for the two genes *norA*, *norC* and *norB* in 11 bacteria (11%) are positive for all three genes (30%).

In a study conducted by Gade et al., resistance to ciprofloxacin and ofloxacin in *S. aureus* isolated from blood infection was reported as 92.5% and 80.4%, respectively (17-18). Another study by Afzal et al. showed 56% of *S. aureus* isolated from eye infection were phenotypically resistant to ciprofloxacin. Also, 21.42% of the resistant isolates have high experssion in *norB* gene (19). The results are slightly different from our research, which is probably due to the different criteria of patients in sample collection.

Hadadi's study showed that among the strains with activated pumps, almost half of them (48%) had increased expression. 57% showed increased expression in the gene of a single pump, usually *norA*, while the remaining 43% isolates showed increased expression in 2 or

more efflux pumps, most of which were norB /C. A recent study of 52 ciprofloxacin-resistant S. aureus bacterial isolates showed that the activity of efflux pumps increased in 23% of cases, which is associated with increased bacterial resistance to fluoroquinolones (20). The results of this research are consistent with the level of gene expression obtained in our research. A plausible explanation for this could be that these isolates had a sufficient number of pumps in their cell wall before the bacterium was exposed to the drug so that there was no need to increase the expression of the relevant genes. Although fluoroquinolones are not used to treat staphylococcal infections, their widespread use in the treatment of nosocomial infections of this bacterium in the hospital has been suggested as the most important factor in the development and spread of S. aureus resistance to fluoroquinolones (20).

In a study conducted in 2016 by Haddadi Zahmatkesh et al. 68% were methicillin and 24% ciprofloxacin-resistant, and the gene frequencies for *norA* and *norB* in resistant strains were 100% and 83%, respectively, which showed that the efflux pump in Ciprofloxacin resistance plays a key role and the presence of these genes can be important in suggesting a therapeutic pattern (20). As in our study, increased expression of light genes was observed in fluoroquinolone-resistant isolates.

Also, Costa et al confirmed the main role of efflux pump as the first agent for antimicrobial resistance in *S. aureus* (5). In 2015, another study showed that all strains of *S. aureus* resistant to ciprofloxacin have the *norA* gene and 83% of the isolates have the *norB* gene (20).

A study in Boston on the expression of *norB* gene and *norC* gene in ciprofloxacin-resistant staphylococci was performed on 115 isolates from 2009 and 2011 isolated from Korea. *norB* was present in 42% of the samples

## Homepages http://jabs.fums.ae.fr



Zargar M, et al.

and *norA* and *norC* genes were present in 33.62% and 53.2% of the samples, respectively (21-22). The difference in the expression of resistance genes in different studies can be due to the difference in the distribution of strains in different geographical locations, depending on the infection control of the location and the amount of resistant isolates.

Based on the role of efflux pumps in bacterial resistance to some antibiotics, studies are being conducted to prevent the expression of each of the efflux pump genes and to evaluate antibiotic resistance. Accordingly, studies by Pourmand et.al in 2014 showed the increased expression of *norA* in methicillin and ciprofloxacin resistant *S. aureus* strain (23).

In our study, the determination of the correlation between the expersion of *nor genes* with fluoroquinolone resistance phenotype is one of the strengths of the study.

#### **Conclusion**

The results of this study showed that pump efflux is one of the important mechanisms of resistance to fluoroquinolone antibiotics such as ciprofloxacin, while the role of other factors involved in resistance should not be ignored. Finally, it is suggested that further studies be performed to produce and extend the molecules of efflux pump inhibitors. It is also recommended that the expression of pump efflux genes in ciprofloxacin-resistant strains be studied by molecular methods such as real-time PCR.

#### Acknowledgements

This study is financially supported by Islamic Azad University, Arak Branch in 2019 code: 97834. The authors of current study would like to express their deep thanks to all laboratory staff of Islamic Azad University, Arak Branch.

#### **Conflict of interests**

The authors declare that they have no competing interests.

### References

- 1.Tseng TT, Gratwick KS, Kollman J, Park D, Nies DH, Goffeau A, et al. The RND permease superfamily: an ancient, ubiquitous and diverse family that includes human disease and development proteins. J Mol Microbiol Biotechnol. 1999;1(1):107–25.
- 2.Levy SB. Active efflux mechanisms for antimicrobial resistance. Antimicrob Agents Chemother. 1992;36(4):695–703.
- 3. Pakzad I, Zayyen Karin M, Taherikalani M, Boustanshenas M, Lari AR. Contribution of AcrAB efflux pump to ciprofloxacin resistance in Klebsiella pneumoniae isolated from burn patients. GMS Hyg Infect Control. 2013;8:2.
- 4.Corredor Arias LF, Luligo Espinal JS, Moncayo Ortiz JI, Santacruz Ibarra JJ, Álvarez Aldana A. Relationship between super antigenicity, antimicrobial resistance and origin of Staphylococcus aureus isolated. Colomb medica Cali, Colomb. 2016;47(1):15–20.
- 5.Costa SS, Junqueira E, Palma C, Viveiros M, Melo-Cristino J, Amaral L, et al. Resistance to Antimicrobials Mediated by Efflux Pumps in Staphylococcus aureus. Antibiot (Basel, Switzerland). 2013;2(1):83–99.
- 6.Costa SS, Sobkowiak B, Parreira R, Edgeworth JD, Viveiros M, Clark TG, et al. Genetic Diversity of norA, Coding for a Main Efflux Pump of Staphylococcus aureus. Front Genet. 2018;9:710.
- 7. Asadi-Samani M, Khaledi M, Khaledi F, Samarghandian S, Gholipour A.Phytochemical properties and antibacterial effects of Salvia multicaulis Vahl., Euphorbia microsciadia Boiss., and Reseda lutea on Staphylococcus aureus and Acinetobacter baumanii. Jundishapur J Nat Pharm Prod 14(3):e63640.
- 8. Andersen JL, He G-X, Kakarla P, KC R, Kumar S, Lakra WS, et al. Multidrug Efflux Pumps from Enterobacteriaceae, Vibrio cholerae and Staphylococcus aureus Bacterial Food Pathogens. International Journal of Environmental Research and Public Health. 2015. 12,p. 1487–547.
- 9. Jenul C, Horswill AR. Regulation of Staphylococcus aureus Virulence. Microbiol Spectr. 2019;7:2.
- 10.Guo Y, Song G, Sun M, Wang J, Wang Y. Prevalence and Therapies of Antibiotic-Resistance in Staphylococcus aureus. Front Cell Infect Microbiol. 2020:10:107.
- 11. Schindler BD, Jacinto P, Kaatz GW. Inhibition of drug efflux pumps in Staphylococcus aureus: current status of potentiating existing antibiotics. Future Microbiol. 2013 Apr;8(4):491–507.
- 12.Dashtbani-Roozbehani A, Brown MH. Efflux Pump Mediated Antimicrobial Resistance by Staphylococci in Health-Related Environments: Challenges and

DOR: 20.1001.1.22285105.2023.13.1.8.9



### Journal of Advanced Biomedical Sciences

## Homepages https://jabs.fums.ac.fr



#### Efflux Pump Genes norA, norB and norC Related to Fluoroquinolones Resistance

- the Quest for Inhibition. Antibiot (Basel, Switzerland). 2021;10(12):12.
- 13.Bostanmaneshrad A, Nowroozi J, Eslami G. Assessing the antibiotic resistance patterns dependent on efflux pump genes of Staphylococcus aureus strains isolated from Blood culture in Shahid Beheshti hospital centers during 1396-1397. Research-in-Medicine. 2021;1;45(2):55–61.
- 14.Karmakar A, Dua P, Ghosh C. Biochemical and Molecular Analysis of Staphylococcus aureus Clinical Isolates from Hospitalized Patients. Can J Infect Dis Med Microbiol J Can des Mal Infect la Microbiol medicale. 2016;2016:9041636.
- 15.Kwiecinski JM, Horswill AR. Staphylococcus aureus bloodstream infections: pathogenesis and regulatory mechanisms. Curr Opin Microbiol. 2020;53:51–60.
- 16.L. JN, Arnold L, H. MM, Weiguo L, R. DM, H. TV, et al. Quinolone Efflux Pumps Play a Central Role in Emergence of Fluoroquinolone Resistance in Streptococcus pneumoniae. Antimicrob Agents Chemother. 2006 1;50(1):310–7.
- 17. Kwak YG, Truong-Bolduc QC, Bin Kim H, Song K-H, Kim ES, Hooper DC. Association of norB overexpression and fluoroquinolone resistance in clinical isolates of Staphylococcus aureus from Korea. J Antimicrob Chemother. 2013;68(12):2766–72.
- 18.Gade ND, Qazi MS. Fluoroquinolone Therapy in

- Staphylococcus aureus Infections: Where Do We Stand? J Lab Physicians. 2013;5(2):109–12.
- 19.Afzal M, Vijay AK, Stapleton F, Willcox M. The Relationship between Ciprofloxacin Resistance and Genotypic Changes in S. aureus Ocular Isolates. Pathog (Basel, Switzerland). 2022;11(11):11.
- 20.Haddadi Zahmatkesh MA, Laripoor M, Mirzaie A, Ashrafi F. Prevalence of norA and norB efflux pump genes in clinical isolates of Staphylococcus aureus and their contribution in ciprofloxacin resistance TT. Iran-J-Med-Microbiol. 2016 1;10(5):20–30. Available from: http://ijmm.ir/article-1-580-en.html
- 21.ruong-Bolduc QC, Strahilevitz J, Hooper DC. NorC, a new efflux pump regulated by MgrA of Staphylococcus aureus. Antimicrob Agents Chemother. 2006; 50(3):1104–7.
- 22.Sun F, Liang H, Kong X, Xie S, Cho H, Deng X, et al. Quorum-sensing agr mediates bacterial oxidation response via an intramolecular disulfide redox switch in the response regulator AgrA. Proc Natl Acad Sci [Internet]. 2012 5;109(23):9095–100. Available from: https://doi.org/10.1073/pnas.1200603109
- 23. Pourmand MR, Yousefi M, Salami SA, Amini M. Evaluation of expression of NorA efflux pump in ciprofloxacin resistant Staphylococcus aureus against hexahydroquinoline derivative by real-time PCR. Acta Med Iran. 2014;52(6):424–9.