



Original Article

## Green Synthesis of Silver Nanoparticles Using *Nasturtium officinale* L. Extract and Analysis of Their Antibacterial Activity Against Nosocomial Pathogens

Shajareh Mahsa Al-Sadat<sup>1</sup>, Dastpak Arezoo<sup>1\*</sup>, Emtiazjoo Mozghan<sup>2</sup>, Mirzaie Amir<sup>3</sup>

1. Department of Biology, Central Tehran Branch, Islamic Azad University, Tehran, Iran

2. Department of Biology, Tehran North Branch, Islamic Azad University, Tehran, Iran

3. Department of Biology, Parand Branch, Islamic Azad University, Parand, Iran

Received: 26 Jan 2022

Accepted: 20 Apr 2022

### Abstract

**Background & Objective:** The green synthesis of the silver nanoparticles (AgNPs) is an eco-friendly and straightforward synthesis method. This study aimed to investigate the green synthesis of AgNPs using *Nasturtium officinale* leaf extract and analysis of their antibacterial activity against some nosocomial pathogens.

**Materials & Methods:** The obtained AgNPs were characterized using UV-visible spectroscopy, X-ray diffraction (XRD), and scanning electron microscope (SEM). In addition, the antibacterial activity of synthesized AgNPs and *N. officinale* leaf extract were performed against five bacterial strains (*Staphylococcus epidermidis*, *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*) by agar disk diffusion assay and minimum inhibitory concentration test.

**Results:** The green synthesized AgNPs had significant antimicrobial activity against all studied bacteria. In addition, greater inhibition activity was observed against *E. coli* and *P. aeruginosa* rather than other bacteria.

**Conclusion:** The synthesized AgNPs revealed a special antibacterial effect against selected bacteria, compared to leaf extract of *N. officinale*. Identifying new antibacterial agents with profound efficacy against antibiotic-resistant opportunistic pathogens is essential. These data provide an eco-friendly and rapid green approach for AgNPs synthesis and the potential of AgNPs for use in drug development against nosocomial infections.

**Keywords:** Nosocomial pathogens, Silver nanoparticles, *Nasturtium officinale*

### Introduction

Nosocomial infections are infections usually acquired after hospitalization and manifest 48-72 hours after admission to the hospital. For nosocomial infections, responsible pathogens include bacteria, fungi, and viruses. In opportunistic bacterial infections, common organisms include

*Staphylococcus epidermidis*, *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* (1, 2). Studies have shown that 5% –10% of hospitalizations in Europe and North America result in nosocomial infections, in Asia and Africa, it increased to 40% (3). Based on National Healthcare Safety Network with the Center for Disease Control (CDC) there are 13 types of nosocomial infection with 50 infection sites (4, 5). Antibiotic-resistant bacteria strains are considered

**\*Corresponding Author:** Dastpak Arezoo, Department of Biology, Central Tehran Branch, Islamic Azad University, Tehran, Iran

Email: arezoo.dastpak@iauctb.ac.ir

<https://orcid.org/0000-0002-7443-3023>

Shajareh Mahsa Al-Sadat: <https://orcid.org/0000-0002-5456-1474>  
Emtiazjoo Mozghan: <https://orcid.org/0000-0002-2654-1951>  
Mirzaie Amir: <https://orcid.org/0000-0002-6569-3413>

a significant threat to health care systems worldwide. Methicillin-resistant *Staphylococcus aureus* (MRSA) is the most common resistant Gram-positive bacterium responsible for serious nosocomial infections leading to death. MRSA is resistant to nearly all beta-lactam antibiotics, due to the expression of penicillin-binding protein 2a (PBP2a). PBP2a is persistent in the action of all Beta-lactam antibiotics. MRSA is now considered a community-related infection (CA-MRSA) due to the prevalence of antibiotic resistance (6, 7). Recently, nanoparticle-based approaches such as nanoliposomes and metal nanoparticles have been developed to overcome these problems. Nanoparticles are made up of dozens or hundreds of atoms or molecules and are found in a wide range of sizes (8, 9). Silver nanoparticles have engrossed a great deal of interest due to potential advantages in various research areas, especially pharmaceutical problems and biomedical applications, such as antimicrobial and anticancer activities (10, 11). Many studies have shown that surface active oxygen species formed on nanosilver are released in certain conditions and may have antimicrobial and antifungal activity (12, 13). Several techniques have been reported for the preparation of nanoparticles. The facet, well-defined shapes, sizes, and controlled synthesis of nanoparticles are crucial in their activity (14, 15).

Various physical and chemical processes have been reported for synthesizing AgNPs such as gamma irradiation, laser ablation, chemical reduction and synthetic biological techniques (16). Due to the growing need for safe, lower toxicity, environmental-friendly

and better size controlling aspects, green synthesis of nanoparticles from various plant extracts and essential oils is the preferred path of nanoparticle preparation over various physical and chemical methods (17). Also, biosynthesis methods do not require high pressure, temperature, energy, or toxic and dangerous chemicals. Besides, plants represent highly stable nanoparticles compared to other techniques and are straightforward to measure (18). Medicinal plants have a great value for detecting medical drugs and people have been using them for various aims from the beginning of the history of humans (19, 20). *Nasturtium officinale* L. (Watercress) is a perennial herb native to Asia and Europe and belongs to the Brassicaceae family that can be grown around and in water (Figure1). The plant is used fresh or as an ingredient in different food (21). It contains vitamins especially A, B, and C, phenolic compounds, folic acid, minerals, fibers, proteins, iron, and calcium, and a high level of glucosinolates which is the highest concentration compared to all other vegetable plants, along with high carotenoids concentration, for example, beta-carotene and lutein (20, 21, 22). Traditionally, the plant has numerous medical uses, such as anti-diabetes, antibacterial, stopping hemorrhages, chest pain, purifying blood, anemia, iron deficiency, throat expectorant, gallbladder, bronchitis, tumors and tuberculosis (23, 24). Previous studies have displayed antioxidant, nephroprotective, anti-inflammatory, hepatoprotective, and anti-hyperlipidemia features of the watercress in vitro and in vivo conditions (20, 23, 24).



Figure 1. *Nasturtium officinale*.



In the present study, a simple and rapid approach was applied for the green synthesis of AgNPs using *Nasturtium officinale* L. leaf extract. Furthermore, the antimicrobial activity of *N. officinale* leaf extracts and nanoparticles synthesized via leaf extracts were evaluated against common nosocomial pathogens.

## Materials & Methods

### Source of chemicals and bacteria

Ethanol, Antibiotics, Dimethylsulfoxide (DMSO), Silver nitrate ( $\text{AgNO}_3$ ) were taken from Sigma-Aldrich (St Louis, MO, USA). Mueller Hinton broth (MHB) was obtained from Oxoid, Basingstoke, United Kingdom. Mueller-Hinton agar (MHA) was purchased from Merck, Germany. All other reagents were of analytical grade and used as received. All strains of bacteria were obtained from the Pasteur Institute of Iran.

### Preparation of plant samples

*Nasturtium officinale* L. (Watercress) was collected from the mountains of Mako city in the West Azerbaijan province, Iran. Botanists identified the plant parts at the biology department, Islamic Azad University. Plant materials were washed thoroughly with distilled water to remove any earthy matter or remove the dirt and any other contaminations. The plant leaves were dried at room temperature under shade to retain their fresh green color and prevent the loss of active compounds. Dried leaves of *N. officinale* were powdered by a grain mill and kept in glass containers for further use.

### Preparation of plant extract

The powder of *N. officinale* leaves was subjected to ethanol extraction whereby 30g of powder leaves were macerated in 300 ml of ethanol 80% for 72h at room temperature. The resulting mixture was filtered (Whatman No. 2) and then evaporated at 33°C using a rotary evaporator with an 80 rpm rotation speed. The stock solution is prepared in 10% dimethylsulfoxide (DMSO) and was stored at 4°C.

### Synthesis of silver nanoparticles

6 ml of the plant extract and 40 ml of 10 mM aqueous silver nitrate solution were mixed with continuous stirring for 5 minutes at room temperature to synthesize silver nanoparticles. The colorless silver nitrate solution turns to brown color that reveals the formation of AgNPs. Silver nanoparticles were collected by centrifugation (13,000 rpm for 20 min) and dried at 40°C for two h, before evaluation of the characterization.

### Characterization of silver nanoparticles UV-visible spectroscopic characterization of AgNPs

Silver ions ( $\text{Ag}^+$ ) reduction to metallic particles ( $\text{Ag}^0$ ) and formation of AgNPs were identified by measuring the UV-Vis spectrum using UV-visible spectrophotometer (JASCO V-670 Spectrophotometer) within the wavelength ranges of 300–700 nm (25).

### X-ray diffraction analysis

The X-ray diffraction (XRD) analyses were recorded for AgNPs to evaluate the crystal and nano-structural of AgNPs. The spectra were obtained on a Philips PW-1730 X-ray diffractometer (XRD) in the range of  $2\theta = 0^\circ$ – $110^\circ$ , using Copper  $K\alpha$  radiation.

### Scanning electron microscopy

The surface characteristics and morphology of the prepared AgNPs were investigated by scanning electron microscopy (SEM) (XL30, Philips, Eindhoven, Holland).

### Bacterial strains

For further use, all organisms were maintained on Mueller Hinton broth (MHB) containing 30% (v/v) glycerol at -20°C for further use. The antibacterial properties of ethanolic extracts of *N. officinale* leaves were evaluated using the following strains of bacteria, Gram-negative bacteria: *Pseudomonas aeruginosa* (ATCC 27853), *Escherichia coli* (ATCC 25922), *Klebsiella pneumoniae* (ATCC 700603),



and Gram-positive bacteria: *Staphylococcus aureus* (ATCC 25923) and *Staphylococcus epidermidis* (ATCC 12228).

### Antibacterial Activity Assay

Bacterial suspensions equivalent to 0.5 McFarland ( $1.5 \times 10^8$  CFU/ml) were made in sterile normal saline solution. A sterile cotton swab was dipped into a test tube containing bacterial suspensions and then was cultured on the Mueller-Hinton agar (MHA). These plates were incubated for 24 h at 37°C.

### Determination of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)

Briefly, The extract and green synthesized nanoparticles MICs, separately, were determined using the broth microdilution assay following recommendations of the CLSI guidelines 2018 (26). The MIC test was carried out using the standard broth microdilution method. It was done in a sterile 96-well round bottom microtitre plate.

A total of 95 µl of MHB and 5 µl of every bacterial strain suspension were added to the 96 wells. 100 µl of plant extract (800 mg/ml DMSO) was then added into the first wells of each column and serial dilution was prepared. A sterility control well and a growth control well were also studied for each strain. The same was done for green synthesized AgNPs (2000 µg/ml MBH). The plate was incubated for 24 h at 37 °C. The last well which had no bacterial growth was defined as the MIC.

MBC values were determined using sub-culturing of bacterial suspension ( $10^8$  CFU/mL) from the MIC tubes into Mueller-Hinton agar plates and then incubated at 37°C for 24h. The diameter of the growth inhibition zone (mm) surrounding the disks for plant extracts, AgNPs and antibiotics showed the inhibitory effect.

### Disk Diffusion Assay

The antibacterial activities of both ethanolic

extract and green silver nanoparticles synthesized were tested for antibacterial properties against bacterial strains and then compared with antibiotics (Tetracycline, Erythromycin, Chloramphenicol, and Ciprofloxacin) using the agar disk diffusion techniques according to Clinical and Laboratory Standards Institute (CLSI) guidelines 2018 (26). They are the most common antibiotics in treating nosocomial infections (27-29). Briefly, each strain was spread on a sterile Mueller-Hinton agar plate using a sterile cotton swab. For testing the antibacterial properties of extract, the sterile 6 mm blank paper disks (Padtan Teb Inc., Tehran, Iran) were loaded with 50 µL of filter-sterilized plant extracts and four disks with antibiotics as the positive control, and the corresponding volume of ethanol was used as the negative control (30, 31). For evaluation of the antimicrobial property of synthesized nanoparticles, the sterile paper disks were impregnated with 50 µL of AgNPs (62.5 µg/ml), and four disks with antibiotics and one disk were as negative control. After 24 h of incubation at 37°C, the inhibitory zone was measured for extract, AgNPs, and antibiotics. For each sample, three replicate trials were conducted against each organism.

### Statistical Analysis


The statistical analyses were conducted using SPSS statistics applying the student's t-test and presented as mean  $\pm$  SD. A p-value less than 0.05 was considered statistically significant.

## Result

### Synthesis of AgNPs

The formation of AgNPs was visually indicated by color change (from light yellow to dark brown) in the mixtures of silver nitrate solution and leaf extract (Figure 2). The confirmation of NPs synthesis was characterized by UV-vis spectroscopy, XRD, and SEM.






**Figure 2.** Visual observation of synthesis of AgNPs. A) *N. officinale* leaf extract, B) *N. officinale* leaf extract treated

### UV-visible Spectroscopy

The UV-visible spectroscopic analysis is one of the valuable and reliable techniques to confirm the formation of nanoparticle synthesis (18). A broad absorption peak at around 421.5 nm confirmed AgNPs were successfully gained by plant extract (Figure 3, A)

### X-ray diffraction analysis

X-ray diffraction (XRD) proved the crystalline nature of silver nanoparticles. The XRD patterns for synthesized AgNPs revealed diffraction peaks at  $2\theta$  degrees = 38.31, 44.43, 64.73, and 77.74 corresponding to (111), (200), (220), and (311) reflections, respectively (Figure 3, B).



**Figure 3A.** UV-visible absorption spectrum of AgNPs synthesized using *N. officinale* leaf extract treated with 10mM  $\text{AgNO}_3$ .

**B.** XRD pattern of AgNPs synthesized by *N. officinale* leaf extract


### Scanning electron microscopy analysis of silver nanoparticles

The morphology and particle size of the AgNPs were evaluated by scanning electron microscopy. This technique magnifies the image by using electrons instead of light. The high-density AgNPs synthesized by treating *N. officinale* extract are observed in the SEM image. The average silver nanoparticle size was 22.64 nm and had a spherical shape (Figure 4, A).


### Antibacterial Activity

### Disk Diffusion Assay

A disk diffusion test was performed to evaluate the sensitivity of standard strains to plant ethanolic extracts and synthesized silver nanoparticles. For assessing the antibacterial activity of extract of *N. officinale* leaves, biosynthesis AgNPs, and antibiotics (Tetracycline, Erythromycin, Chloramphenicol, and Ciprofloxacin) on bacteria strains, diameters of inhibition growth zones were measured and the results are presented as shown in Table 1, Figure 4, B. and Chart 1.



**Figure 4.** A SEM micrograph showed the morphology of synthesized AgNPs using *N. officinale* leaf extract. B. Antibacterial activities of silver nanoparticles *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Staphylococcus aureus* and *Staphylococcus epidermidis* compared with antibiotics (Tetracycline, Erythromycin, Chloramphenicol, and Ciprofloxacin)



**Chart 1.** Comparative antibacterial analysis of *Nasturtium officinale* L. extracts with some common antibiotics based on the disk diffusion method

The disc diffusion assay showed *N. officinale* extract has antibacterial activities against all the tested bacteria. The highest activity of the plant extract was found against Gram-negative bacteria *P. aeruginosa* ( $11 \pm 0.37$  mm) and *E. coli* ( $10 \pm 0.22$  mm). The growth inhibition was moderately active against bacteria *S. aureus* ( $9 \pm 0.34$  mm), *K. pneumoniae* ( $9 \pm 0.23$  mm) and *S. epidermidis* ( $8 \pm 0.16$  mm).

The in vitro test for antibacterial activity revealed that AgNPs synthesis by *N. officinale* leaf extract inhibited all of the growth of Gram-negative and Gram-positive, with an inhibition zone of  $13 \pm 0.36$  mm to  $16 \pm 0.32$  mm. The positive control (Tetracycline, Erythromycin, Chloramphenicol, Ciprofloxacin) showed different inhibition zone depending on Bacteria strains (Table 1).

**Table 1.** Inhibition zone (IZ) of leaf extract of *Nasturtium officinale* and biosynthesized silver nanoparticles (AgNPs) against five pathogenic bacteria

Bacteria	Antibiotics	IZ of antibiotics(mm) (mean $\pm$ SD)	IZ of AgNPs (mm) (mean $\pm$ SD)	IZ of extract (mm) (mean $\pm$ SD)
<i>Escherichia coli</i>	TE	22 $\pm$ 0.23	13 $\pm$ 0.36	10 $\pm$ 0.22
	ER	8 $\pm$ 0.31		
	CH	24 $\pm$ 0.44		
	CI	40 $\pm$ 0.19		
<i>Staphylococcus epidermidis</i>	TE	18 $\pm$ 0.41	14 $\pm$ 0.25	8 $\pm$ 0.16
	ER	12 $\pm$ 0.18		
	CH	23 $\pm$ 0.43		
	CI	30 $\pm$ 0.31		



<i>Staphylococcus aureus</i>	TE	20 ± 0.24	15 ± 0.47	9 ± 0.34
	ER	10 ± 0.41		
	CH	28 ± 0.47		
	CI	34 ± 0.34		
<i>Klebsiella pneumoniae</i>	TE	28 ± 0.32	15 ± 0.25	9 ± 0.23
	ER	11 ± 0.37		
	CH	31 ± 0.23		
	CI	33 ± 0.25		
<i>Pseudomonas aeruginosa</i>	TE	20 ± 0.21	16 ± 0.32	11 ± 0.37
	ER	10 ± 0.26		
	CH	17 ± 0.36		
	CI	39 ± 0.26		

\*TE: Tetracycline, ER: Erythromycin, CH: Chloramphenicol. CI: Ciprofloxacin

### MIC and MBC of ethanolic extract and silver nanoparticles

MIC and MBC of each strain were specified for both biosynthesis AgNPs and *N. officinale* extract. MIC and MBC of the plant extracts were assessed against all the experimented bacteria (Table 2). To compare the effect of the plant extract and green synthesis nanoparticles,

the MIC and MBC were remarkable. Table 3 showed that AgNPs exhibited the lowest MIC and MBC against *S. aureus* (62.5µg/ml), and plant extract had the lowest value for MIC against *S. aureus* and *S. epidermidis* (25mg/ml). Also, the highest MBC was related to *E. coli* with a concentration of 250 µg/ml for biosynthesis AgNPs.

**Table 2.** MIC and MBC (mg/ml) values of *Nasturtium officinale* ethanolic extract against studied bacteria

Bacteria	MIC(mg/ml)	MBC(mg/ml)
<i>Escherichia coli</i>	50	200
<i>Klebsiella pneumoniae</i>	50	100
<i>Pseudomonas aeruginosa</i>	50	100
<i>Staphylococcus aureus</i>	25	100
<i>Staphylococcus epidermidis</i>	25	200



**Table 3.** MIC and MBC ( $\mu\text{g/ml}$ ) of the AgNPs against selected bacteria

Bacteria	MIC(mg/ml)	MBC(mg/ml)
<i>Escherichia coli</i>	125	250
<i>Klebsiella pneumoniae</i>	125	125
<i>Pseudomonas aeruginosa</i>	125	125
<i>Staphylococcus aureus</i>	62.5	62.5
<i>Staphylococcus epidermidis</i>	125	125

## Discussion

Recently, the increase in opportunistic bacterial infections and the development of antibiotic-resistant infections associated with high mortality are becoming worrisome. Therefore, detecting novel antibacterial agents against opportunistic pathogens is urgently required (32). Advances in nanobiotechnology, novel applications of nanoparticles, and research on natural products toward detecting affective antibacterial agents have become extensive attention to the synthesis of nanoparticles using plants (18, 16). The chemical synthesis of AgNPs requires the use of harmful chemicals that may affect the environment and human health. The synthesis of nanoparticles using plants offers several advantages. It presents the best manner to synthesize silver nanoparticles due to the numerous active biomolecules such as alkaloids, phenolic acids, polyphenols, terpenoids found in plants (33, 34).

This study assayed the silver nanoparticles produced by *N. officinale* leaf extract and their antibacterial effects. AgNPs from *N. officinale* were successfully synthesized. Silver nanoparticles display a brown color in an aqueous solution due to a surface plasmon excitation in AgNPs that agrees with previous studies (35, 36). Similar color changes have also been observed in different studies, confirming the chemical reaction between leaf extract and silver nitrate. Also, UV–Vis spectroscopy, SEM, and XRD analysis were carried out to characterize the nanoparticles. Following previous studies, the maximum absorption peak in the UV-Vis spectra around 421.5 nm confirms the formation of Ag nanoparticles (36, 37). The XRD pattern revealed the purity of synthesized AgNPs without additional diffraction peaks. The crystalline structure of synthesized AgNPs was



characterized through X-Ray comparison to standards. The spherical shape of silver nanoparticles was displayed by SEM image. The bioactive compounds found in plant extracts can be helpful for reducing metal ions and resulting in the synthesis of silver nanoparticles (38, 39). The SEM image displayed that the size of AgNPs synthesized was 16 - 35 nm. A similar result of the size of the biosynthesized silver nanoparticles by *N. officinale* extract has been reported in Sadeghi, (2014) study (40). The disk diffusion test and MIC / MBC assays were performed against both Gram-positive and Gram-negative bacteria to assess the potential of green nanosilver bactericidal effects. It should be attended that the antibacterial properties of AgNPs depend on their structure including the size, shape, colloidal state, preparation method, and the bacteria with which they interact (37, 38, 41). The exact mechanisms by which nanosilver performs antibacterial efficacy are still under verification. It was considered that AgNPs form pits on the bacterial cell wall surface leading to the demolition of the cell structures and damaging cell function (42-44). Besides, it seems ionic Adhesion of the AgNPs onto the bacteria cell wall and plasma membrane, owing to the interaction between positively charged Ag ions and negative surface charge of the cell membrane, causes structural variations in the membrane and the disorder of proton motive force and thus destruction of membrane. Moreover, silver nanoparticles by manufacturing free radicals upon contact with bacteria harm the membrane (45-48). The large sizes of zones growth inhibition produced by *N. officinale* extract, AgNps, and the four antibiotics (Tetracycline, Erythromycin, Chloramphenicol, and Ciprofloxacin) against all bacteria indicated the potency of the active constituents in *N. officinale*, silver nanoparticles, and antibiotics. Ethanolic extract of *N. officinale* revealed the highest zone of inhibition in *P. aeruginosa* and *E. coli*. The Gram-negative bacteria have extra tolerant against a broad range of antibiotics.

The presence of lipopolysaccharides, mutations in porins and changing the hydrophobic properties in the outer membrane can cause this resistance (49-51). The extract gives a robust inhibitory effect against Gram-negative bacteria; this may demonstrate the more function of *N. officinale* extract specifically against the Gram-negative cell membrane and cell wall.

Synthesized AgNPs are effective against Gram-negative and Gram-positive bacteria with the highest inhibitory zone in *P. aeruginosa* (Chart 1). Some studies have shown that the formed AgNPs using plant extract were effective against negative and positive pathogens. Silver nanoparticles synthesis by *N. officinale* had appreciable antibacterial activity and the higher concentration was essential for growth inhibition of *E. coli*. (52-54). Compared to the antibiotics, the plant extract was more effective than Erythromycin on *P. aeruginosa* and *E. coli*. In contrast, synthesized AgNPs demonstrated a greater effect on all five bacteria strains than Erythromycin. Ciprofloxacin and Chloramphenicol had the greatest and lowest effect on all bacteria. The lowest MIC value was achieved for *S. aureus* exhibiting that both extract and AgNPs have the maximum toxicity to this bacteria (25mg/mL and 62.5 µg/mL, respectively). *S. epidermidis* is another sensitive Gram-positive microorganism to plant extract its MIC value was 25mg/ml. The results indicated AgNPs have considerably higher antibacterial activity than plant extract and 62.5 µg/ml was sufficient to inhibit *S. aureus* growth. Different amounts in MIC of extracts may be due to various chemical components and the volatile temper of the constituents (55). Several studies have shown the inhibitory effects of synthesized AgNPs on the Staphylococcus aureus, which was consistent with the present study's findings (56, 57). Mahdavi et al. (2019) showed that *E. coli* had a high resistance to *N. officinale* essential oil as this study revealed resistance *E. coli* to leaf extract (58). Also, another study revealed that natural hexanol in the plant extracts,



including *N. officinale*, affected bacteria such as *S. aureus* and *E. coli* (59). This is one of the components in *N. officinale* extract that causes inhibiting the growth of bacteria.

### Conclusion

The presence of adverse microorganisms in human health has become a significant concern, due to the variety of infections and diseases. Thus, there is a need for better effective and nontoxic treatment options with more antimicrobial activity against infectious diseases, incredibly opportunistic bacterial infections and resistant strains. Plant-based silver nanoparticles (AgNPs) have specified antimicrobial properties that deem them fit for use as an alternative to antibiotics and have many advantages like cost-effectiveness, efficiency, safety and less toxicity to living organisms. The antimicrobial properties shown by the AgNPs reported here represent a great alternative to achieving this objective. The synthesis reported here provides a green and eco-friendly method to rapidly synthesize AgNPs through implementing *Nasturtium officinale* leaf extract at ambient temperature. The study's findings propose that the leaf extract of *N. officinale* is an appropriate reducing and capping agent for the biosynthesis of AgNPs. Initially, the observation of changing the color of Ag<sup>+</sup> solution from colorless to dark brown confirmed the formation of AgNPs. The synthesized AgNPs were characterized using UV-Visible, XRD and SEM analysis. In addition, the antibacterial properties of both ethanolic extract and nanoparticles obtained by extract were assessed against five tested bacteria by disc-diffusion and MIC/MBC methods. The results suggest the synthesized AgNPs are more effective than extract and low concentrations are required to kill both Gram-negative and Gram-positive bacteria. Thus, this rapid, environmentally friendly, and cost-effective method and significant antibacterial activity may potentially suggest using such nanoparticles in future drug development and the food industry. Nevertheless, research in pharmacological and toxicological studies, especially in vivo, is

needed to develop and design future antimicrobial therapeutic agents for in vivo application.

### Conflicts of interest

We declare that we have no conflict of interest.

### Acknowledgment

The authors acknowledge support provided for this project by Central Tehran Branch, Islamic Azad University, Tehran, Iran. (Thesis code: 10160205932008)

### References

1. Hemeg HA. Nanomaterials for alternative antibacterial therapy. *Int J Nanomedicine*. 2017; 12:8211-8225.
2. Monegro AF, Muppidi V, Regunath H. Hospital-acquired infections. StatPearls. Available from: <https://www.coursehero.com/file/67681574/Discussiondocx.2020>.
3. McFee RB. Nosocomial or hospital-acquired infections: an overview. *Dis Mon*. 2009; 55(7):422-438.
4. Khan HA, Ahmad A, Mehboob R. Nosocomial infections and their control strategies. *Asian Pac J Trop Biomed*. 2015; 5(7):509-514.
5. Sharma S, Shabir S. The Menace of nosocomial infections. *Glob J. Oto*. 2017;7(4):555717.
6. Kateete DP, Bwanga F, Seni J, Mayanja R, Kigozi E, Mujuni B, et al. CA-MRSA and HA-MRSA coexist in community and hospital settings in Uganda. *Antimicrob Resist Infect Control*. 2019; 8(1):1-9.
7. Siddiqui AH, Koirala J. Methicillin Resistant *Staphylococcus aureus*. [Updated 2021 Jul 19]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK482221/>
8. Eid KA, Azzazy HM. Sustained broad-spectrum antibacterial effects of nanoliposomes loaded with silver nanoparticles. *Nanomedicine*. 2014; 9(9):1301-1310.
9. Roy I, Ohulchanskyy TY, Pudavar HE, Bergey EJ, Oseroff AR, Morgan J, et al. Ceramic-based nanoparticles entrapping water-insoluble photosensitizing anticancer drugs: a novel drug-carrier system for photodynamic therapy. *J Am Chem Soc*. 2003; 125(26):7860-7865.
10. Savithramma N, Rao ML, Rukmini K, Suvarnalatha Dp. Antimicrobial activity of silver nanoparticles synthesized by using medicinal plants. *Int J Chemtech Res*. 2011; 3(3):1394-1402.
11. Tran QH, Le AT. Silver nanoparticles: synthesis, properties, toxicology, applications and perspectives. *Adv Nat Sci Nanosci Nanotechnol*. 2013; 4(3):033001.
12. Levard C, Hotze EM, Lowry GV, Brown GE Jr. Environmental transformations of silver nanoparticles: impact on stability and toxicity. *Environ Sci Technol*. 2012; 46(13):6900-6914.



13. Prabhu S, Poulose EK. Silver nanoparticles: mechanism of antimicrobial action, synthesis, medical applications, and toxicity effects. *Int Nano Lett.* 2012; 2(1):1-10.
14. Khan I, Saeed K, Khan I. Nanoparticles: Properties, applications and toxicities. *Arab J Chem.* 2019; 12(7):908-931.
15. Pugazhenthiran N, Anandan S, Kathiravan G, Prakash NKU, Crawford S, Ashokkumar M. Microbial synthesis of silver nanoparticles by *Bacillus sp.* *J Nanopart Res.* 2009; 11(7):1811-1815.
16. Iravani S, Korbekandi H, Mirmohammadi SV, Zolfaghari B. Synthesis of silver nanoparticles: chemical, physical and biological methods. *Res Pharm Sci.* 2014; 9(6):385-406.
17. Bhardwaj B, Singh P, Kumar A, Kumar S, Budwar V. Eco-friendly greener synthesis of nanoparticles. *Adv Pharm Bull.* 2020; 10(4):566-576.
18. Devatha CP, Thalla AK. Green synthesis of nanomaterials. Synthesis of inorganic nanomaterials. UK: Woodhead Publishing. 2018.
19. Farnsworth NR. The role of ethnopharmacology in drug development. *Ciba Found Symp.* 1990; 154:2-11.
20. Faizy HS, Esmail LS, Mahdi HS. Phytochemicals Analysis in Watercress (*Nasturtium officinale*) Plant Extracts, IOP Conference Series: Earth and Environmental Science. 2021; 761: 012042
21. Hassandokht M, Jafari S, Ebrahimi R. Watercress (*Nasturtium officinale* R. Br.) Breeding. In: Al-Khayri J.M., Jain S.M., Johnson D.V. (eds) *Advances in Plant Breeding Strategies: Vegetable Crops*. Springer. Cham. Available from: [https://doi.org/10.1007/978-3-030-66969-0\\_6](https://doi.org/10.1007/978-3-030-66969-0_6). 2021.
22. O'Neill ME, Carroll Y, Corridan B, Olmedilla B, Granado F, Blanco I, et al. A European carotenoid database to assess carotenoid intakes and its use in a five-country comparative study. *Br. J. Nutr.* 2001; 85(4):499-507
23. Alqahtani MA, Al Othman MR, Mohammed AE. Bio fabrication of silver nanoparticles with antibacterial and cytotoxic abilities using lichens. *Sci Rep.* 2020; 10(1):16781
24. Sedaghattalab M, Razazan M, Sadeghi H, Hossein Doustimotlagh A, Akbartabar Toori M, Abbasi Larki R, et al. Effects of *Nasturtium officinale* Extract on Antioxidant and Biochemical Parameters in Hemodialysis Patients: A Randomized Double-Blind Clinical Trial. *Evid Based Complementary Altern. Med.* 2021; 2021: 1-8.
25. Kumari J, Baunthiyal M, Singh A. Characterization of silver nanoparticles synthesized using *Urtica dioica* leaves and their synergistic effects with antibiotics. *J Radiat Res Appl Sci.* 2016;9(3):217-227.
26. Weinstein MP, Patel JB, Burnham CA, Campeau S, Conville PS, Doern C, et al. Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial disk susceptibility tests. CLSI Standard M02, Clinical and Laboratory Standards Institute. 13th ed. USA;2018.
27. Eliakim-Raz N, Lador A, Leibovici-Weissman Y, Elbaz M, Paul M, Leibovici L. Efficacy and safety of chloramphenicol: joining the revival of old antibiotics? Systematic review and meta-analysis of randomized controlled trials. *J Antimicrob Chemother.* 2015;70(4):979-996.
28. Hansen S, Sohr D, Piening B, Pena Diaz L, Gropmann A, Leistner R, et al. Antibiotic usage in German hospitals: results of the second national prevalence study. *J Antimicrob Chemother.* 2013; 68(12):2934-2939.
29. Hassan MM. Scenario of Antibiotic Resistance in Developing Countries. In: Mareş, M, Lim S, Lai K, Cristina R, editors. *Antimicrobial Resistance - A One Health Perspective* [Internet]. London: IntechOpen; 2020. Available from: <https://www.intechopen.com/chapters/74593> doi: 10.5772/intechopen.94957
30. Azarmehr N, Afshar P, Moradi M, Sadeghi H, Alipoor B, Khalvati B, et al. Hepatoprotective and antioxidant activity of watercress extract on acetaminophen-induced hepatotoxicity in rats, *Heliyon.* 2019; 5(7): e02072.
- 31- Moyo M, Gomba M, Nharingo T. Afzelia quanzensis bark extract for green synthesis of silver nanoparticles and study of their antibacterial activity. *Int J Ind Chem.* 2015;6:329-338.
32. Frieri M, Kumar K, Boutin A. Antibiotic resistance. *J Infect Public Health.* 2017; 10(4):369-378.
33. Das VL, Thomas R, Varghese RT, Soniya EV, Mathew J, Radhakrishnan EK. Extracellular synthesis of silver nanoparticles by the *Bacillus* strain CS 11 isolated from industrialized area. *3 Biotech.* 2014;4(2):121-126.
34. He Y, Du Z, Lv H, Jia Q, Tang Z, Zheng X, et al. Green synthesis of silver nanoparticles by *Chrysanthemum morifolium* Ramat. extract and their application in clinical ultrasound gel. *Int J Nanomedicine.* 2013; 8:1809-1815.
35. Amendola V, Bakr OM, Stellacci F. A Study of the surface plasmon resonance of silver nanoparticles by the discrete dipole approximation method: Effect of shape, size, structure, and assembly. *Plasmonics.* 2010; 5:85-97.
36. Praba PS, Vasantha, V, Jeyasundari J, Jacob YBA. Synthesis of plant-mediated silver nanoparticles using *Ficus microcarpa* leaf extract and evaluation of their antibacterial activities. *Eur Chem Bull.* 2015; 4(1):117-120.
37. Akintelu SA, Bo Y, Folorunso AS. A Review on synthesis, optimization, mechanism, characterization, and antibacterial application of silver nanoparticles synthesized from plants. *J Chem.* 2020;2020:1-12.
38. Benakashani F, Allafchian AR, Jalali SAH. Biosynthesis of silver nanoparticles using *Capparis spinosa* L. leaf extract and their antibacterial activity. *Karbala Int J Mod Sci.* 2016; 2(4):251-258.
39. Jemal K, Sandeep BV, Pola S. Synthesis, characterization, and evaluation of the antibacterial activity of *Allophylus serratus* leaf and leaf derived callus extracts mediated silver nanoparticles. *J Nanomater.* 2017; 2017: 1-11.





40. Sadeghi B. Synthesis of silver nanoparticles using leaves aqueous extract of *Nasturtium officinale* (NO) and its antibacterial activity. *Int J Mol Clin Microbio*. 2014; 4(2):428-434.
41. Gonzalez-Fernandez S, Lozano-Iturbe V, Garcia B, Andres LJ, Menendez MF, Rodriguez D, et al. Antibacterial effect of silver nanorings, *BMC. Microbiol*. 2020; 20(1):172-187
42. Ayala-Núñez NV, Villegas HHL, Turrent L, Padilla CR. Silver nanoparticles toxicity and bactericidal effect against Methicillin-Resistant *Staphylococcus aureus*: Nanoscale does matter. *Nanobiotechnol.*, 2009; 5:2-9.
43. Korshed P, Li L, Liu Z, Wang T. The Molecular mechanisms of the antibacterial effect of picosecond laser generated silver nanoparticles and their toxicity to human cells. *PLoS One*. 2016; 11(8):0160078.
44. Nalwade AR, Jadhav A. Biosynthesis of silver nanoparticles using leaf extract of *Datura alba* Nees. and evaluation of their antibacterial activity. *Arch Appl Sci Res*. 2013; 5(3): 45-49.
45. Kędziora A, Speruda M, Krzyżewska E, Rybka J, Łukowiak A, Bugla-Płoskońska G. Similarities and differences between silver ions and silver in nanoforms as antibacterial agents. *Int J Mol Sci*. 2018; 19(2):444.
46. Kvítek L, Panáček A, Soukupová J, Kolář M, Večeřová R, Prucek R, et al. Effect of surfactants and polymers on stability and antibacterial activity of silver nanoparticles (NPs). *J Phys Chem*. 2008; 112(15):5825-5834.
47. Mikhailova EO. Silver Nanoparticles: Mechanism of Action and Probable Bio-Application, *J. Funct. Biomater*. 2020; 11(4):84.
48. Tamboli DP, Lee DS. Mechanistic antimicrobial approach of extracellularly synthesized silver nanoparticles against gram positive and gram negative bacteria. *J Hazard Mater*. 2013; 260:878-884.
49. Breijyeh Z, Jubeh B, Karaman R. Resistance of gram-Negative bacteria to current antibacterial agents and approaches to resolve it. *Molecules*. 2020; 25(6):1340.
50. Sheldon A. Antibiotic mechanisms of action and resistance. In: Textbook of diagnostic microbiology, In: Mahon CR, Lehman DC, Manuselis G. editors. 3rd ed. St. Louis: Saunders ;2007.
51. Razmavar S, Abdulla MA, Ismail SB, Hassandarvish P. Antibacterial activity of leaf extracts of *Baeckea frutescens* against methicillin-resistant *Staphylococcus aureus*. *Biomed Res Int*. 2014; 2014:521287.
52. Awwad AM, Salem NM, Aqarbeh MM, Abdulaziz FM. Green synthesis, characterization of silver sulfide nanoparticles and antibacterial activity evaluation. *Chem. Int*. 2020; 6(1):42-48.
53. Gupta A, Saleh NM, Das R, Landis RF, Bigdeli A, Motamedchaboki K, et al. Synergistic antimicrobial therapy using nanoparticles and antibiotics for the treatment of multidrug-resistant bacterial infection. *IOP science.*, 2017; 1(1): 15004.
54. Lee JH, Lim JM, Velmurugan P, Park YJ, Park YJ, Bang KS, Oh BT. Photobiologic-mediated fabrication of silver nanoparticles with antibacterial activity. *J Photochem Photobiol B*. 2016;162:93-99.
55. Gonelimali FD, Lin J, Miao W, Xuan J, Charles F, Chen M, et al. Antimicrobial properties and mechanism of action of some plant extracts against food pathogens and spoilage microorganisms. *Front Microbiol*. 2018; 24(9):1639.
56. Chaloupka K, Malam Y, Seifalian AM. Nanosilver as a new generation of nanoparticle in biomedical applications. *Trends biotechnol*. 2010;28(11):580-588.
57. Sadeghi BA. Green synthesis of silver nanoparticles using seed aqueous extract of *Olea europaea*. *Int J Nanodimens*. 2014; 5(6):575-581.
58. Mahdavi S, Kheyrollahi M, Sheikhlouei H, Isazadeh A. Antibacterial and antioxidant activities of essential oil on food borne bacteria', *Open. Microbiol. J*. 2019; 13(1):81-85.
59. Cha JD, Jeong MR, Jeong SI, Moon SE, Kim JY, Kil BS, et al. Chemical composition and antimicrobial activity of the essential oils of *Artemisia scoparia* and *A. capillaris*. *Planta Med*. 2005; 71(2):186-190.